



**North West
School of Psychiatry**

Semester 3 Handbook

MRCPsych Course

2020 – 2022

A Psychiatry Medical Education Collaborative between mental health Trusts and Health Education North West.

Course Director - Dr Latha Hackett, Consultant in Child & Adolescent Psychiatry

Deputy Course Director - Dr Dushyanthan Mahadevan, Consultant in Child & Adolescent Psychiatry

List of Contributors

Course Director

Dr Latha Hackett, Consultant in Child and Adolescent Psychiatry

Latha.hackett@mft.nhs.uk

Deputy Course Director

Dr Dushyanthan Mahadevan, Consultant in Child & Adolescent Psychiatry

Dushyanthan.Mahadevan@elht.nhs.uk

Module Leads

Across the Ages	Dr Latha Hackett	Latha.hackett@mft.nhs.uk
CAMHS	Dr Neelo Aslam	Neelo.aslam@mft.nhs.uk
Forensic	Dr Victoria Sullivan & Dr Amit Sharda	Victoria.sullivan@gmmh.nhs.uk Amit.Sharda@lancashirecare.nhs.uk
General Adult	Dr Sally Wheeler & Dr Swanand Patwardhan	sally.wheeler@gmmh.nhs.uk Swanand.Patwardhan@gmmh.nhs.uk
Intellectual Disabilities	Dr Sol Mustafa	sol.mustafa@nhs.net
Old Age	Dr Anthony Peter	Anthony.peter@lancashirecare.nhs.uk
Psychotherapy	Dr Adam Dierckx	Adam.Dierckx@gmmh.nhs.uk
Substance Misuse	Dr Patrick Horgan	Patrick.Horgan2@gmmh.nhs.uk

Trust Leads

CWP	Dr Matthew Cahill	matthew.cahill1@nhs.net
GMMH (NMGH)	Dr Swanand Patwardhan	Swanand.Patwardhan@gmmh.nhs.uk
GMMH (Prestwich)	Dr Catrin Evans & Dr Emily Mountain	Catrin.Evans@gmmh.nhs.uk Emily.Mountain@gmmh.nhs.uk
Lancashire Care	Dr Clare Oakley	Clare.Oakley@lancashirecare.nhs.uk
Mersey Care	Dr Indira Vinjamuri (covering for Dr Yenal Dundar)	indira.vinjamuri@merseycare.nhs.uk
NWBH	Dr Yogesh Sharma	Yogesh.Sharma@nwbh.nhs.uk
Pennine Care	Dr Ema Etuk	ema.etuk@nhs.net

Medical Education Managers (MEM's)

CWP	Jonathan Ruffler	jonathan.ruffler@nhs.net
GMMH	Sam Abbott	sam.abbott@gmmh.nhs.uk
Lancashire Care	Jacqueline Welding	jacqueline.welding@lancashirecare.nhs.uk
Mersey Care	Dawn McLoughlin	dawn.mcloughlin@merseycare.nhs.uk
NWBH	Ian Bithell	ian.bithell@nwbh.nhs.uk
Pennine Care	Kimberley McDowell	kimberley.mcdowell@nhs.net

Contents

Brief guidelines for case conference presentation.....	9
Brief guidelines for journal club presentation.....	10
Syllabus Links.....	11
Curriculum Mapping.....	12
Links to Critical Appraisal Checklists.....	13
General Adult.....	14
Session 13: Psychosis-3	14
Learning Objectives	14
Expert Led Session.....	14
Case Presentation	14
Journal Club Presentation (Select 1 paper).....	14
'555' Topics (Select 1 topic; 5 slides with no more than 5 bullet points)	14
Session 14: Depression-3	15
Learning Objectives	15
Expert Led Session.....	16
Case Presentation	16
Journal Club Presentation (Select 1 paper).....	16

'555' Topics (Select 1 topic; 5 slides with no more than 5 bullet points)	16
MCQs	16
Session 15: Bipolar Disorder-3	17
Learning Objectives	17
Expert Led Session.....	18
Case Presentation	18
Journal Club Presentation (Select 1 paper).....	18
'555' Topics (Select 1 topic; 5 slides with no more than 5 bullet points)	18
MCQs	19
Session 16: Anxiety disorders-2 (GAD, panic disorder, phobic anxiety disorders)	20
Learning Objectives	20
Expert Led Session.....	20
Case Presentation	21
Journal Club Presentation (Select 1 paper).....	21
'555' Topics (Select 1 topic; 5 slides with no more than 5 bullet points)	21
MCQs	21
Session 17: Suicide/self-harm-2	23
Learning Objectives	23
Expert Led Session.....	23
Case Presentation	23
Journal Club Presentation (Select 1 paper).....	23
'555' Topics (Select 1 topic; 5 slides with no more than 5 bullet points)	23
MCQs	23
Session 18: Perinatal psychiatry	25
Learning Objectives	25
Expert Led Session.....	25
Case Presentation	25

Journal Club Presentation (Select 1 paper).....	25
'555' Topics (Select 1 topic; 5 slides with no more than 5 bullet points)	25
MCQs	26
CAMHS	28
Session 5: Attachment	28
Learning Objectives.....	30
Curriculum Links.....	30
Expert Led Session	30
Case Presentation	30
Journal Club Presentation	30
'555' Topics (1 slide on each topic with no more than 5 bullet points).....	31
MCQs	31
Additional Resources /Reading Materials.....	34
Session 6: Assessment of Mental Health Problems in Child & Adolescents with Intellectual Disability (ID)	34
Learning Objectives	34
Curriculum Links	36
Expert Led Session.....	36
Case Presentation	36
Journal Club Presentation	36
'555' Topics (1 slide on each topic with no more than 5 bullet points)	36
MCQs	37
Additional Resources /Reading Materials.....	40
Old Age	39
Session 5: Mood Disorders in the Older Person	39
Learning Objectives	39
Curriculum Links.....	39
Expert Led Session.....	39

Case Presentation	39
Journal Club Presentation.....	39
'555' Topic (5 slides with no more than 5 bullet points per slide).....	40
MCQs	40
Additional Resources / Reading Materials	41
Session 6: Psychosis in the Older Person	47
Learning Objectives	47
Curriculum Links	47
Expert Led Session.....	47
Case Presentation	47
Journal Club Presentation.....	47
'555' Topic (5 slides with no more than 5 bullet points per slide).....	47
MCQs	48
Additional Resources / Reading Material	49
MCQ answers	51
Mood disorders	51
Psychosis	51
Across the Ages	52
Coming Soon	
Forensic	53
Session 3: Too mad to murder?	53
Learning Objectives.....	53
Curriculum Links	53
Expert Led Session	53
Case Presentation.....	54
Journal Club Presentation	54

'555' Topic (5 slides with no more than 5 bullet points per slide)	55
MCQs.....	55
Additional Resources / Reading Materials	57
ID	60
Session 3: Behavioural Issues in Intellectual Disability	60
Learning Objectives	60
Curriculum Links	60
Expert Led Session.....	60
Case Presentation	60
Journal Club Presentation.....	60
'555' Topics (5 slides on each topic with no more than 5 bullet points)	61
MCQ's	61
Additional Resources/ Reading Materials	62
Psychotherapy	63
Session 3: Psychological approaches to Depression	63
Learning Objectives	63
Curriculum Links	63
1.1, 1.2, 1.3, 1.3.4, 2.3, 2.4, 2.6, 2.8, 6.1, 7.1.1, 9, 14.....	63
Expert Led Session.....	63
Case Presentation	63
Journal Club Presentation.....	63
'555' Topics (5 slides on each topic with no more than 5 bullet points)	63
MCQs	64
Additional Resources / Reading Materials	65
Substance Misuse	66

Session 3: Diagnosis and Management of People with co-occurring mental health and alcohol/drug use conditions 66

Learning Objectives 66

Curriculum Links 66

Expert Led Session..... 66

Case Presentation 66

Journal Club Presentation..... 66

'555' Topics (5 slides on each topic with no more than 5 bullet points) 67

MCQs 67

MCQ Answers 69

Additional Resources / Reading Materials 70

Brief guidelines for case conference presentation

The objectives of case conference are:

- 1) To provide a forum to discuss complex/interesting cases in a learning atmosphere.
- 2) To develop your ability to present cases in a concise and logical manner.
- 3) To develop your presentation skills.

Guidelines for presenters:

1. Please use PowerPoint for the presentation (or if you are using other tools make sure that they are compatible with your local IT facilities).
2. You have to present a case that is relevant to the theme of the day on which you are presenting.
3. Please meet with your educational/clinical supervisor at least 4-6 weeks prior to the presentation to identify an appropriate case to present. If there is no suitable case in the team that you work in, you may have to approach other teams/consultants to identify a case.
4. Cases can be chosen for their atypical presentation, diagnosis, complexity or for exploring management options.
5. It would be helpful if you can identify specific clinical questions that you would like to be discussed/answered at the end of the presentation.
6. We would recommend the following structure for the presentation:
 - Introduction (include reasons for choosing the case)
 - Circumstances leading to admission (if appropriate)
 - History of presenting complaint
 - Past Psychiatric history
 - Medical History/ current medication
 - Personal/family History
 - Alcohol/Illicit drugs history
 - Forensic history
 - Premorbid personality
 - Social circumstances
 - Mental state examination
 - Investigations
 - Progress since admission (if appropriate)

- A slide with questions that you would you like to be discussed
- Discussion on differential diagnosis including reasons for and against them.
- Management / treatment

7. The structure of the presentation can vary as long it is logical and concise. Please build into the presentation some natural points to stop and discuss the case.

8. Important: Please ask a senior member of your team who knows the case to attend on the day you are presenting.

Brief guidelines for journal club presentation

The objectives of journal club presentation are:

- 1) To learn to perform a structured critical appraisal of a study.
- 2) To learn to make appropriate use of evidence in making decisions about the care of your patients.
- 3) To prepare for the MRCPsych exams.
- 4) To develop your presentation skills.

Guidelines for presenters:

1. Please use PowerPoint for the presentation (or if you are using other tools make sure that they are compatible with your local IT facilities).
2. Please select one of the 3 papers listed for the week from the School of Psychiatry handbook to present.
3. Email the paper to your local co-ordinator at least a week before the presentation so that it can be circulated in time.
4. As the presenter you are expected to both present the paper and critically review it.
5. We would recommend the following structure for the presentation: Background to study, methods, analysis, results, conclusions, critical appraisal of the study and implications for clinical practice
6. The most important part of the presentation is the critical appraisal. This should include aspects such as:
 - Purpose of the study

- Type of study
- Subject selection and any bias
- Power calculation (could the study ever answer the question posed)
- Appropriateness of statistical tests used
- Use of relevant outcomes
- Implications of findings
- Applications of findings/conclusions in your area
- Directions for further research

7. Use standardized critical appraisal tools.

8. Please discuss the paper and the presentation with your educational/clinical supervisor prior to the presentation.

Syllabus Links

[Syllabus for MRCPsych](#)

[Syllabus for MRCPsych critical review](#)

MRCPsych [Paper A](#) -The Scientific and theoretical basis of Psychiatry

MRCPsych [Paper B](#) - Critical review and the clinical topics in Psychiatry

MRCPsych [CASC](#)

Curriculum Mapping				
Section	Topic	Covered by		
		LEP	AP	LR
7.1	Disorders in adulthood			
7.1.1	Unipolar depression	✓		✓
7.1.2	Bipolar depression	✓		✓
7.1.3	Schizophrenia	✓		✓
7.1.4	Anxiety disorders	✓		✓
7.1.5	OCD	✓		✓
7.1.6	Hypochondriasis		✓	✓
7.1.7	Somatization disorder		✓	✓
7.1.8	Dissociative disorders		✓	✓
7.1.9	Personality disorders	✓		✓
7.1.10	Organic psychoses	✓		✓
7.1.11	Other psychiatric disorders	✓		✓
7.2	Perinatal Psychiatry		✓	✓
7.3	General Hospital Psychiatry		✓	✓
7.4	Emergency Psychiatry*		✓	✓
7.5	Eating Disorders			
7.5.1	Anorexia nervosa		✓	✓
7.5.2	Bulimia nervosa		✓	✓
7.6	Psycho-sexual disorders			
7.6.1	Non-organic sexual dysfunction, etc.		✓	✓
7.6.2	Gender Identity Disorders		✓	✓
-	Mental Health Act 1983	✓		✓

Key- LEP – Local Education Programme;

AP- Academic Programme

LR – Learning Resources

Links to Critical Appraisal Checklists	
Study	Checklists
Randomized Controlled Trial	<ol style="list-style-type: none"> 1. CONSORT Checklist 2. SIGN Checklist 3. CASP Checklist
Case-control Study	<ol style="list-style-type: none"> 1. SIGN Checklist 2. CASP Checklist
Cohort Study	<ol style="list-style-type: none"> 1. SIGN Checklist 2. CASP Checklist
Meta-analysis & Systematic Review	<ol style="list-style-type: none"> 1. PRISMA statement 2. SIGN Checklist 3. CASP Checklist
Qualitative study	<ol style="list-style-type: none"> 1. CASP Checklist
Economic study	<ol style="list-style-type: none"> 1. SIGN Checklist 2. CASP Checklist
Diagnostic study	<ol style="list-style-type: none"> 1. SIGN Checklist 2. CASP Checklist

General Adult
Session 13: Psychosis-3
Journal theme: Meta-analysis / Systematic Review on Psychosis
Learning Objectives
<ul style="list-style-type: none"> • To develop an understanding of the biopsychosocial management of schizophrenia • To develop an understanding of evidence-based treatment • To develop an understanding of the use of antipsychotics in special cases e.g. liver and renal impairment • To develop an understanding of Meta-analysis / Systematic Review and develop skills for critically appraising them.
Expert Led Session
<ul style="list-style-type: none"> • Schizophrenia: Biopsychosocial management and evidence based treatment.
Case Presentation
<ul style="list-style-type: none"> • A case of Schizophrenia (any subtype) /Schizoaffective disorder / Delusional disorder / Acute and transient psychotic disorder / First-episode psychosis
Journal Club Presentation (Select 1 paper)
<ul style="list-style-type: none"> • Leucht S, Komossa K, Rummel-Kluge C, Corves C, Hunger H, Schmid F, Asenjo Lobos C, Schwarz S, Davis JM (2009). A meta-analysis of head-to-head comparisons of second-generation antipsychotics in the treatment of schizophrenia. Am J Psychiatry, 166(2):152-63. doi: 10.1176/appi.ajp.2008.08030368 • Souza JS, Kayo M, Tassell I, Martins CB, & Elkisa H. (2013). Efficacy of olanzapine in comparison with clozapine for treatment-resistant schizophrenia: evidence from a systematic review and meta-analyses. CNS Spectrums; 18 (2), 82- 89. DOI: http://dx.doi.org/10.1017/S1092852912000806 • Leucht S, Cipriani A, Spineli L, Mavridis D, Örey D. (2013). Comparative efficacy and tolerability of 15 antipsychotic drugs in schizophrenia: a multiple-treatments meta-analysis. The Lancet; 382 (9896), 951–962. DOI: http://dx.doi.org/10.1016/S0140-6736(13)60733-3
'555' Topics (Select 1 topic; 5 slides with no more than 5 bullet points)
<ul style="list-style-type: none"> • Recommendations for antipsychotics in liver disease • Recommendations for antipsychotics in renal impairment • Antipsychotics and sexual side effects
Statistics '555' topic
<ul style="list-style-type: none"> • t tests, p values and statistical significance

MCQs

1. Which one of the following led a trial that proved Clozapine's effectiveness in treating resistant schizophrenia?
 - A. Kretschmer
 - B. Cade
 - C. Kraepelin
 - D. Kane
 - E. Bleurer
2. Choose the correct match from the following pairs:
 - A. Risperidone: dibenzoxapine
 - B. Droperidol: butyrophenones
 - C. Aripiprazole: benzisothiazole
 - D. Thioridazine: diphenyl butyl piperidine
 - E. Flupentixol: dihydroindole
3. Which of the following atypical agents have the shortest half-life?
 - A. Quetiapine
 - B. Aripiprazole
 - C. Olanzapine
 - D. Clozapine
 - E. Risperidone
4. The patients who are prescribed clozapine or olanzapine should have their serum lipids measured every:
 - A. 6 days whilst on treatment
 - B. One year whilst on treatment
 - C. 3 months for the first year of treatment
 - D. 6 weeks for the first year of treatment
 - E. 6 months for the first year of treatment
5. What percentage of patients develop Tardive Dyskinesia with every year of typical antipsychotic exposure?
 - A. More than 50%
 - B. 2-5%
 - C. 5-10%
 - D. 20-25%
 - E. 10-20%

Session 14: Depression-3

Journal theme: Qualitative study on depression

Learning Objectives

- To develop an understanding of the biopsychosocial management of Depression.
- To develop an understanding of evidence based treatment.
- To develop an understanding of the use of antidepressant in special cases e.g. liver and renal impairment.

- To develop an understanding of Qualitative study and develop skills for critically appraising them.

Expert Led Session

- Depression- Biopsychosocial management and evidence-based treatment

Case Presentation

- A case of major depressive disorder / severe depression with psychotic symptoms / dysthymia / recurrent depressive disorder

Journal Club Presentation (Select 1 paper)

- Coupe N, Anderson E, Gask L, Sykes P, Richards DA, et al. (2014). Facilitating professional liaison in collaborative care for depression in UK primary care; a qualitative study utilising normalisation process theory. BMC Family Practice; 15:78. DOI: 10.1186/1471-2296-15-78
- Gask L, Rogers A, Oliver D, May C, Roland M (2003) [Qualitative study of patients' perceptions of the quality of care for depression in general practice](#). Br J Gen Pract., 53(489):278-83.
- Gask L, Dixon C, May C, Dowrick C (2005) [Qualitative study of an educational intervention for GPs in the assessment and management of depression](#). Br J Gen Pract, 55(520):854-9.

'555' Topics (Select 1 topic; 5 slides with no more than 5 bullet points)

- ECT – indications and contraindications
- Depression – important rating scales
- Treatment of refractory depression- first choice (evidence-based)

Statistics '555' topic

- Basic statistics- mean, median, mode, range, standard deviation, standard error

MCQs

1. Which of the following neurotransmitters does Venlafaxine act on?
 - A. Serotonin only
 - B. Noradrenaline and Serotonin
 - C. Dopamine
 - D. Noradrenaline, Serotonin and Dopamine
 - E. GABA

2. Which of the following statements about Trazodone is FALSE?
- It is relatively safe in overdose
 - It does not have strong antihistamine properties
 - It is not a MAO-A and MAO- B inhibitor
 - It does not block 5-HT reuptake
 - It is a 5HT2 agonist
3. Which of the following are not common side effects of Sertraline?
- Tachycardia and arrhythmias
 - Nausea, vomiting, abdominal pain
 - Sexual dysfunction
 - Agitation, anxiety
 - Insomnia
4. Laura is a depressed 61-year-old woman who has not responded to an SSRI and has urinary incontinence. Which one of the following antidepressants is the best choice in this situation?
- Phenelzine
 - Mirtazapine
 - Vortioxetine
 - Trazodone
 - Duloxetine
5. Hypertension is a common side effect of which of the following antidepressants?
- Venlafaxine
 - Paroxetine
 - Escitalopram
 - Trazodone
 - Mirtazapine

Session 15: Bipolar Disorder-3
Journal theme: RCT on bipolar disorder
Learning Objectives

- To develop an understanding of the biopsychosocial management of Bipolar disorder.
- To develop an understanding of evidence-based treatment.
- To develop an understanding of the use of mood-stabilizers in special cases e.g. liver and renal impairment.

- To develop an understanding of Randomized Controlled trials and develop skills for critically appraising them.

Expert Led Session

- Bipolar disorder- Biopsychosocial management and evidence-based treatment.

Case Presentation

- A case of type I bipolar disorder / type II bipolar disorder / cyclothymia / bipolar disorder with psychotic symptoms / rapid cycling bipolar disorder/ unipolar mania.

Journal Club Presentation (Select 1 paper)

- Jones SH, Smith G, Mulligan LD, Lobban F, Law H, et al. (2010). Recovery-focused cognitive-behavioural therapy for recent-onset bipolar disorder: randomised controlled pilot trial. *The British Journal of Psychiatry*; 206 (1) 58-66. DOI: 10.1192/bjp.bp.113.141259
- Castle D, White C, Chamberlain J, Berk M, Berk L, Lauder S, Murray G, Schweitzer I, Piterman L, Gilbert M (2010) [Group-based psychosocial intervention for bipolar disorder: randomized controlled trial](#). *BJPsych*, 196: 383-388.
- Kemp D, Gao K, Fein E, Chan P, Conroy C, Obral S, Ganocy S, Calabrese R (2012) [Lamotrigine as add-on treatment to lithium and divalproex: lessons learned from a double-blind, placebo-controlled trial in rapid-cycling bipolar disorder](#). *Bipolar Disord.*, 14(7):780-789.

'555' Topics (Select 1 topic; 5 slides with no more than 5 bullet points)

- Monitoring requirements for mood stabilizers
- Bipolar I Vs Bipolar II – differences in diagnosis and treatment
- Role of Psychotherapy in long term management of Bipolar Disorder

Statistics '555' topic

- Intention-to-treat analysis & Last Observation Carried Forward (LOCF)

MCQs

1. Sodium valproate:
 - A. Is mostly renally metabolised
 - B. Commonly causes hypertrichosis
 - C. Reduces lamotrigine levels
 - D. Is licensed for prophylaxis of BPAD
 - E. Is a first line choice in acute mania
2. Which of the following drugs has a high therapeutic index:
 - A. Lithium
 - B. Carbamazepine
 - C. Phenytoin
 - D. Warfarin
 - E. Gabapentin
3. The risk of Ebstein's anomaly in babies born to woman taking lithium is:
 - A. 1:10
 - B. 1:100
 - C. 1:500
 - D. 1:1000
 - E. 1:10000
4. Which of the following commonly causes hypercalcaemia:
 - A. Lithium
 - B. Valproate
 - C. Risperidone
 - D. Quetiapine
 - E. Clozapine
5. Lithium levels in once daily nocte dosing should be taken:
 - A. 4 hours post dose
 - B. 12 hours post dose
 - C. 6 hours post dose
 - D. Immediately before the next dose
 - E. 8 hours post dose

6 . Match the following to a drug from the list below:

1. Spina Bifida
 2. Tricuspid valve defect
 3. Cleft palate
 4. Microcephaly
- A - Lithium
B - Benzodiazepines
C - Valproate
D - None of the above

7. Match the following mood stabilisers to their chemical structure:

1. Haloperidol D
 2. Risperidone A
 3. Olanzapine C
 4. Quetiapine B
- A. Benzizoxazole
B. Dibenzothiazepine
C. Thienobenzodiazepine
D. Butyrophenone

Session 16: Anxiety disorders-2 (GAD, panic disorder, phobic anxiety disorders)

Journal theme: case –control studies on the topic

Learning Objectives

- To develop an understanding of GAD, panic disorder, phobic anxiety disorders (aetiology, epidemiology, natural history, neurobiology, genetics, diagnostic criteria, classification, psychopathology, clinical presentation, assessment, risks) and their management (pharmacological, psychological, social).
- To develop an understanding of Case-control studies and develop skills for critically appraising them.

Expert Led Session

- Biopsychosocial management of GAD, panic disorder and phobic anxiety disorders.

Case Presentation

- A case where either GAD, panic disorder or phobic disorder is the main diagnosis or a differential diagnosis.

Journal Club Presentation (Select 1 paper)

- Lipka J, Miltner WH, Straube T (2011) [Vigilance for threat interacts with amygdala responses to subliminal threat cues in specific phobia](#). Biol Psychiatry, 70(5):472-8.
- Santos MA, Ceretta LB, Réus GZ, Abelaira HM, Jornada LK, Schwalm MT, Neotti MV, Tomazzi CD, Gulbis KG, Ceretta RA, Quevedo J (2014) Anxiety disorders are associated with quality of life impairment in patients with insulin-dependent type 2 diabetes: a case-control study. Rev Bras Psiquiatr., 36 (4):298-304.
- Kiropoulos L, Klien B, Austin D, Gilson K, Pier C, Mitchell J and Ciechowski L (2008) [Is internet-based CBT for panic disorder and agoraphobia as effective as face-to-face CBT?](#) Journal of anxiety disorders 22(8), 1273-1284.

'555' Topics (Select 1 topic; 5 slides with no more than 5 bullet points)

- CBT for agoraphobia- principles
- Principles of use of benzodiazepines for anxiety disorders
- Prognosis in GAD and Panic Disorder

Statistics '555' topic

- Null hypothesis, Type-1 error and type-2 error

MCQs

1. Venlafaxine is not licenced for which of the following indications?
 - A. Social anxiety
 - B. PTSD
 - C. Panic disorder
 - D. Depression +/- Anxiety
 - E. GAD
2. The following are TRUE of the pharmacokinetics of benzodiazepines:
 - A. When long-acting they have long elimination half-life.

- B. When short-acting they have a small distribution volume.
- C. When long-acting they have no active metabolites
- D. When short-acting they have high accumulation
- E. Benzodiazepines with a half-life of 12 hours tend to be used as anxiolytics.

3. Which of the following statements is FALSE about the effects of hypnotics on sleep?

- A. Benzodiazepines suppress stage IV sleep.
- B. With chronic Benzodiazepines use suppression of REM sleep in the early part of the night occurs
- C. On withdrawal of Benzodiazepines a rebound increase above the 'normal' amount of REM sleep occurs.
- D. It may take up to 6 weeks to see a return to a normal sleep pattern on Benzodiazepine withdrawal.
- E. Barbiturates are more likely to suppress REM sleep than are Benzodiazepines.

4. With regards to the NICE guidelines on GAD, which of the following is FALSE?

- A. SSRIs (particularly Sertraline) are the first line medications.
- B. SNRIs are second line.
- C. If the patient cannot tolerate SSRI or SNRI, offer Pregabalin.
- D. Antipsychotic should be offered for the treatment of GAD in primary care.
- E. Do not offer a benzodiazepine for the treatment of GAD in primary or secondary care except as a short-term measure during crises

5. With respect to the NICE guidelines on psychological intervention for GAD, which of the following is FALSE?

- A. CBT for people with GAD should be based on the treatment manuals used in the clinical trials of CBT for GAD.
- B. CBT for GAD usually consist of 12–15 weekly sessions (fewer if the person recovers sooner; more if clinically required), each lasting 1 hour.
- C. Practitioners providing high-intensity psychological interventions for GAD need not have regular supervision to monitor fidelity to the treatment model.
- D. If a person with GAD chooses a high-intensity psychological intervention, offer either CBT or applied relaxation.
- E. Consider providing all interventions in the preferred language of the person with GAD if possible.

Session 17: Suicide/self-harm-2 Journal theme: Any study method on the topic
Learning Objectives
<ul style="list-style-type: none"> To develop an understanding of various facets of self-harm and suicide (aetiology, epidemiology, neurobiology, genetics, clinical presentation, risk assessment) and their management (pharmacological, psychological, social).
Expert Led Session
<ul style="list-style-type: none"> Suicide & self-harm- comprehensive risk assessment
Case Presentation
<ul style="list-style-type: none"> Cases related to any type of clinical presentations where suicide and/ or self-harm is the central theme
Journal Club Presentation (Select 1 paper)
<ul style="list-style-type: none"> Quinlivan L, Cooper J, Steeg S, Davies L, Hawton K, Gunnell D, Kapur N (2014) Scales for predicting risk following self-harm: an observational study in 32 hospitals in England. BMJ Open, doi: 10.1136/bmjopen-2013-004732. Kapur N, Gunnell D, Hawton K, Nadeem S, Khalil S, Longson D, Jordan R, Donaldson I, Emsley R, Cooper J (2013) Messages from Manchester: pilot randomised controlled trial following self-harm. BJPsych 203: 73-74. Kapur N, Turnbull P, Hawton K, Simkin S, Sutton L, Mackway-Jones K, Bennewith O, Gunnell D (2005) Self-poisoning suicides in England: a multi-centre study. Q J Med, 98: 589-597.
‘555’ Topics (Select 1 topic; 5 slides with no more than 5 bullet points)
<ul style="list-style-type: none"> National Confidential Inquiry into suicide by people with mental illness – Key findings of the latest annual report Purpose of a Coroner’s Inquest Epidemiological factors associated with suicides
Statistics ‘555’ topic
<ul style="list-style-type: none"> Standard error and confidence intervals
MCQs
<p>1. Which of the following has been shown to be associated with increased rates of suicide?</p> <p>A. Peptic ulcer</p> <p>B. Non-delusional body dysmorphia</p>

- C. Huntington's chorea
- D. Epilepsy
- E. All of the above

2. Deliberate self-harm is more common in:

- A. Males
- B. Rural areas
- C. Age over 35 years
- D. Lower socioeconomic status
- E. Married

3. Predictors of repetition of DSH include all except:

- A. Personality disorder
- B. Alcohol misuse
- C. Male gender
- D. Previous DSH
- E. Younger age of onset

4. Which of the following is associated with suicide in patients with schizophrenia?

- A. Akathisia
- B. Older patient
- C. Poor premorbid functioning
- D. Short duration of illness
- E. Presence of positive symptoms

5. Which of the following physical health problems are associated with increased risk of suicide?

- A. AIDS
- B. Huntington's disease
- C. Cushing's disease
- D. Porphyria
- E. All of the above

<p>Session 18: Perinatal psychiatry</p> <p>Journal theme: Study with any method</p>
<p>Learning Objectives</p>
<ul style="list-style-type: none"> To understand the impact / risks of major mental disorders on pregnancy and post-partum period. To understand the general principles of prescribing; and the risks & benefits of prescribing psychotropic medications in pregnancy, post-partum period and breast feeding.
<p>Expert Led Session</p>
<ul style="list-style-type: none"> Evidence-based recommendations for psychotropic medications in pregnancy [antipsychotics, antidepressants, mood stabilizers and anxiolytics].
<p>Case Presentation</p>
<ul style="list-style-type: none"> A case of any mental disorder in pregnancy or post-partum period.
<p>Journal Club Presentation (Select 1 paper)</p>
<ul style="list-style-type: none"> Ennis, Z. and Damkier, P. (2015). Pregnancy Exposure to Olanzapine, Quetiapine, Risperidone, Aripiprazole and Risk of Congenital Malformations. A Systematic Review. <i>Basic & Clinical Pharmacology & Toxicology</i>, 116(4), pp.315-320. Boden, R., Lundgren, M., Brandt, L., Reutfors, J., Andersen, M. and Kieler, H. (2012). Risks of adverse pregnancy and birth outcomes in women treated or not treated with mood stabilisers for bipolar disorder: population based cohort study. <i>BMJ</i>, 345(nov07 6), pp.e7085. Uguz, F. (2016). Second-Generation Antipsychotics During the Lactation Period: A Comparative Systematic Review on Infant Safety. <i>Journal of Clinical Psychopharmacology</i>, 36(3), pp.244-252.
<p>'555' Topics (Select 1 topic; 5 slides with no more than 5 bullet points)</p>
<ul style="list-style-type: none"> Post-partum risks of relapse in schizophrenia, bipolar disorder and depression Congenital malformations associated with Lithium, Valproate, Carbamazepine, Lamotrigine and Paroxetine- salient points Use of SSRIs in pregnancy – salient points
<p>Statistics '555' topic</p>
<ul style="list-style-type: none"> Absolute risk reduction, relative risk reduction, Number needed to treat and number needed to harm

MCQs

1. During pregnancy the following physiological changes occur
 - A. Plasma volume markedly increases and eGFR increases
 - B. Plasma volume markedly decreases and eGFR increases
 - C. Plasma volume markedly increases and eGFR decreases
 - D. Plasma volume markedly decreases and eGFR decreases
 - E. There is no change in either plasma volume or eGFR
2. Which of the following is NOT associated with exposure to SSRIs in the Perinatal period?
 - A. Perinatal Death
 - B. Persistent Pulmonary Hypertension of the Newborn
 - C. Postpartum haemorrhage
 - D. Poor neonatal adaptation syndrome
 - E. Preterm birth
3. Which of the following statements is TRUE regarding NICE guidelines?
 - A. Benzodiazepines can be offered in pregnancy for medium term treatment of anxiety
 - B. Consideration of medication dose changes do not have to be made during pregnancy
 - C. If this is a first pregnancy a women's previous response to medication should not influence the choice of antidepressant (being pregnant dictates the choice)
 - D. Lithium can be continued if the women is at high risk of relapse and an antipsychotic is unlikely to be effective
 - E. Measure prolactin levels in women planning pregnancy who are taking a prolactin raising antipsychotic as raised prolactin increases the chances of conception
4. Which of the following statements is TRUE?
 - A. Valproate is associated with reduced fertility in women and men
 - B. Taking Folic acid 5mg with Valproate will reduce teratogenicity
 - C. Valproate monotherapy is not associated with an increased risk of Attention Deficit Hyperactivity Disorder

- D. Valproate monotherapy only affects the child in the 1st and 3rd trimester
- E. Valproate passes in higher concentrations than Lamotrigine in breastmilk

5. Which of the following is TRUE regarding breastfeeding?

- A. Patients with postpartum mental health disorders who require pharmacotherapy should generally be discouraged from breastfeeding
- B. All psychotropic medications are transferred to breast milk in varying amounts
- C. Psychotropics should be chosen with regard to longer half life and less protein binding
- D. Mothers should change their pregnancy medication for breastfeeding
- E. Methadone and Nicotine Replacement Therapy are incompatible with breastfeeding

CAMHS Session 5: Attachment

Learning Objectives

- Describe the concept of attachment and its relevance for the mental health of children and young people.
- To understand the relevance of attachment theory to emotional development, affect regulation and relationships across the lifespan.
- To understand the different classifications of attachment, the conditions that promote healthy attachment or otherwise and the clinical relevance of failure to develop selective attachments.

Curriculum Links

Attachment disorders:

10.8.1.1 10.8.1.2 10.8.1.3 10.8.1.4 10.8.1.5

Expert Led Session

- Should cover assessment, diagnostic challenges and MDT approach in managing attachment disorder. Can also discuss the role of specialist LAC services.

Case Presentation

- To discuss key features in history and presentation and discuss overlap with intrinsic disorders, such as ASD/ADHD.

Journal Club Presentation

- Quasi-autistic patterns following severe early global privation. English and Romanian Adoptees (ERA) Study Team. Rutter M et al. J Child Psychol Psychiatry. 1999 May; 40(4): 537-49.
- Specificity and heterogeneity in children's responses to profound institutional privation. Rutter ML1, Kreppner JM, O'Connor TG; English and Romanian Adoptees (ERA) study team. Br J Psychiatry. 2001 Aug; 179:97-103.

- Genetic, environmental and gender influences on attachment disorder behaviours. Minnis H, Reekie J, Young D, O'Connor T, Ronald A, Gray A, Plomin R. Br J Psychiatry. 2007 Jun; 190:490-5.
- The impact of a statewide trauma-informed care initiative in child welfare on the well-being of children and youth with complex trauma. Bartlett, JD et al. Children and Youth Services Review. 2018 84(1), pp.110-117

'555' Topics (1 slide on each topic with no more than 5 bullet points)

- Evidence based interventions in attachment disorder
- Risk factors for attachment disorder
- Comorbid diagnosis in attachment disorder

MCQs

1. The biological basis of attachment behaviour is:
- A. The child developing relationships with other children
 - B. The mother wanting to protect her child from any harm
 - C. The child seeking proximity to the attachment figure
 - D. The mother's instinct to rear children
 - E. All of the above

2. Attachment theory has been developed by:

- A. Freud
- B. Bowlby
- C. Skinner
- D. Piaget
- E. Klein

3. Fearfulness and “frozen watchfulness” are part of which ICD 10 diagnosis:

- A. Generalised anxiety disorder
- B. Phobic anxiety disorder
- C. PTSD
- D. Reactive attachment disorder
- E. Paranoid personality disorder

4. Select a feature that does NOT form part of Reactive Attachment Disorder (ICD 10) but points towards Pervasive Developmental Disorders:

- A. Abnormal pattern of social responsiveness that improves if child is placed in normal rearing environment
- B. Aggressive responses towards their own or other’s distress
- C. Restricted, repetitive interests and behaviours
- D. Strongly contradictory social responses
- E. None of the above

5. Reactive Attachment Disorder of early infancy and childhood (DSM V) and Reactive Attachment Disorder (ICD 11) share common diagnostic criteria. Which of the following is NOT a diagnostic feature in ICD 11:

- A. Developed before age of 5 years
- B. Does not turn to primary caregiver for comfort
- C. Does not display security seeking behaviour
- D. Does not respond when comfort is offered
- E. None of the above

6. Which of the following features is NOT part of Disinhibited Social Engagement Disorder (ICD 11):

- A. Approaches adults indiscriminately
- B. Exhibits overfamiliar behaviour with strangers
- C. Features develop within the first 5 years of life
- D. Occurs in the context of grossly inadequate childcare
- E. Abnormal speech development including echolalia

7. Which of the following cognitive age ranges must a child reach to develop an attachment relationship:

- A. 2-5 months
- B. 7-9 months
- C. 2 years
- D. 5 years
- E. 7 years

8. What is the procedure called that assesses a child's attachment behaviour:

- A. Novel Situation Test
- B. Attachment Assessment Procedure
- C. Strange Situation Procedure
- D. Mother - Infant Attachment Battery
- E. None of the above

9. Symptoms of Reactive Attachment Disorder have to be present before which age:

- A. 3 years
- B. 9 months
- C. 18 months
- D. 8 years
- E. 5 years

10. The current hypothesis is that Attachment Disorders develop as a result of:

- A. Children having been brought up by a single parent
- B. Children having had limited opportunities to form selected attachments
- C. Children having received a vegetarian diet
- D. Children having intrinsic difficulties in forming secure attachments
- E. Children having a specific gene mutation

Additional Resources / Reading Materials

Books

- Attachment – Attachment and Loss. Bowlby J (first published 1969)
- A Short Introduction to Attachment and attachment disorder – Colby Pearce (second edition 2017)
- Why Love Matters – Sue Gerhardt

E-Learning

- Attachment and how it relates to psychiatry (podcast)

Dr Helen Minnis discusses the issue of attachment in psychiatry and the importance of attunement in the caregiving relationship, taking a look at the current controversies over child care and giving guidance for psychiatrists on how to work with attachment difficulties.

<http://www.psychiatrycpd.org/default.aspx?page=3301>

Growing an Emotional brain: www.youtube.com/watch?v=fzn9OuBqKYs

An introductory blog on Adverse Childhood Experiences (ACEs)

<https://www.connectedforlife.co.uk/blog/2017/6/17/the-adverse-childhood-experiences-ace-study>

And a summary of ACEs literature from a Scottish charity, with hyperlinks to a range of publications and resources, and an educational perspective

<https://www.iriss.org.uk/resources/esss-outlines/aces>

Journal Articles

- The relationship between Adult Attachment and Mental Health Care utilization; a systematic review. Adams G, Wrath A, Meng X. Canadian Journal of Psychiatry 2018 Oct; 63(10): 651–660. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6187440/>
- Making and Breaking of Affectional Bonds. Bowlby J BJPsych 1977 130: 201-10 and 421-431.

Guidelines

NICE guidance on Children's Attachment (2015) <https://www.nice.org.uk/guidance/ng26>

Session 6: Assessment of Mental Health Problems in Child & Adolescents with Intellectual Disability (ID)

Learning Objectives

- To understand the influence of developmental factors on the presentation and treatment of psychiatric disorders.
- To understand the principles and practice of assessment, diagnosis and treatment, including therapeutic modalities, psychoactive medication and environmental manipulations of patients presenting with intellectual disability

Curriculum Links

Intellectual Disability:

13.1 13.2.1 13.2.2 13.3

Expert Led Session

- Should cover assessment and the role of other professionals (OT, LD nurses, LD psychologist) and specialist schools. Evidence based management strategies.

Case Presentation

- To cover presentation and assessment of mental health problems of a child or young person with ID; including how these differ from the non ID population and management strategies (environmental, psychological and medical).
- Those trainees who are not currently in a CAMHS post should contact their local CAMHS team for the suitable case for presentation. Specifically you should identify which Consultants see Children with Learning Disabilities, so an appropriate case can be identified well in advance

Journal Club Presentation

- Vereenooghe L, Flynn S, Hastings RP, et al. Interventions for mental health problems in children and adults with severe intellectual disabilities: a systematic review. *BMJ Open*. 2018;8(6):e021911. Published 2018 Jun 19. doi:10.1136/bmjopen-2018-021911
- Einfeld SL, Ellis LA, Emerson E (2011) Comorbidity of intellectual disability and mental disorder in children and adolescents: A systematic review. *Journal of Intellectual and Developmental Disability* 36 (2) pp137-143.
- Chadwick et al, (2008). Factors associated with the risk of behaviour problems in adolescents with severe intellectual disabilities. *Journal of Intellectual Disability research* 52, (10),864- 876.

'555' Topics (1 slide on each topic with no more than 5 bullet points)

- Cognitive assessment tools in ID (child)
- Child Development assessment tools in paediatrics
- Approaches to assessment in children and young people with limited communication

MCQs

1. People with intellectual disability have previously been classified as:

- A. Mentally retarded
- B. Learning disabled
- C. Sub-normals
- D. Imbeciles
- E. All of the above

2. Intellectual disabilities are defined by which 3 core criteria?

- A. Lower intellectual ability
- B. Onset during childhood
- C. Onset before the age of 8
- D. Significant impairment of social or adaptive functioning
- E. IQ scores are not fixed throughout life

3. Which of the following are generally accepted ranges (ICD-10, DSM-IV) for severity of ID (choose 4)?

- A. Mild (IQ 50-70)
- B. Mild (IQ 70-90)
- C. Moderate (IQ 50-70)
- D. Moderate (IQ 35-50)
- E. Severe (IQ 20-35)
- F. Severe (IQ 25-50)
- G. Profound (IQ below 25)
- H. Profound (IQ below 20)

4. Which of the following 2 statements are true?

- A. Mild ID accounts for approximately 80% of children with ID.
- B. Approximately 50% of children with ID have moderate severity.
- C. Severe ID accounts for approximately 7% of the ID group.
- D. Profound ID affects 10% of children with ID.

5. The prevalence and incidence of ID varies according to gender, age, ethnicity and socioeconomic circumstances. Which statement is false?

- A. Studies generally report a female predominance in LD
- B. Increased maternal age is likely to lead to an increase in incidence of LD
- C. Ethnicity influences prevalence and incidence levels in ID due to the associated links with poverty, access to healthcare, and communications barriers amongst other factors
- D. Lower socioeconomic position is associated with higher prevalence of mild and moderate LD, but not severe LD.

6. Psychiatric illnesses frequently exist comorbidly with ID. Which of the following statements is false?

- A. Prevalence of psychiatric co-morbidity ranges from 30-70%
- B. There is often over diagnosis of co-morbid psychiatric conditions
- C. Practically all categories of mental illness are represented in the ID population
- D. Co-morbid psychiatric problems can vary and change with age

7. Match the following co-morbid problems with the age group they are most likely to present in:

- | | |
|-------------------------------|------------------------|
| 1. Eating and sleep disorders | A. Adolescents |
| 2. Self-injury | B. Very young children |
| 3. ADHD | C. School age children |

8. Which one of the following psychiatric conditions is not generally associated with LD?

- A. Attention deficit hyperactivity disorder
- B. Mood disorders
- C. Anxiety disorders
- D. Psychotic illness
- E. Obsessive compulsive disorder
- F. Anorexia nervosa
- G. Autistic spectrum disorder

9. Behavioural analysis involves which ABC?

- A. Antecedents
- B. Awareness
- C. Boundaries
- D. Behaviour
- E. Consequences
- F. Circumstances

10. Which statement about management of ID is inaccurate?

- A. Medications are commonly under-prescribed when managing challenging behaviour associated with ID.
- B. Behavioural techniques are useful in managing ID
- C. Families provide the majority of support for most people with ID
- D. Social services provide the majority of support for people with ID outside of families

Additional Resources / Reading Materials

Books

- Rutter's Child and Adolescent Psychiatry, Fifth Edition.
Sir Michael Rutter , Dorothy Bishop, Daniel Pine, Steven Scott , Jim S. Stevenson, Eric A. Taylor, Anita Thapar
- Child and Adolescent Psychiatry.
Robert Goodman and Stephen Scott. Third Edition, Wiley-Blackwell

Cerebra resources

<http://w3.cerebra.org.uk/research/research-papers/self-injurious-behaviour-in-children-with-intellectual-disability/>

Journal Articles

- Developing mental health services for Children and Adolescents with Learning Disability: A toolkit for clinicians
<http://www.rcpsych.ac.uk/pdf/devmhservcaldbk.pdf>
- Mental health of children with learning disabilities. Pru Allington-Smith, Advances in Psychiatric Treatment, 2006, vol. 12, 130–140.

Nice Guidelines

Challenging behaviour and learning disabilities: prevention and interventions for people with learning disabilities whose behaviour challenges

<https://www.nice.org.uk/guidance/ng11>

Old Age

Session 5: Mood Disorders in the Older Person

Learning Objectives

- The overall aim of the sessions is for the trainees to gain an overview of mood disorders in later life.
- By the end of the session trainees should:
 - Understand the epidemiology, aetiology and the classification of mood disorders in the elderly.
 - Understand how mood disorders present in the elderly
 - Understand the assessment process including rating scales and neuroimaging
 - Understand the principles of treatment, including treatment resistance.
 - Understand the increased risk of suicide in the elderly.

Curriculum Links

- Old Age Section of the MRCPsych Curriculum: 8.3, 8.4, 8.5, 8.6, 8.7, 8.8, 8.9, 8.10.

Expert Led Session

- A Consultant led session based on the learning objectives listed above.

Case Presentation

- A case to be presented which highlights key issues in an older person presenting with a mood disorder. Please consider the learning objectives above.

Journal Club Presentation

- Hedna, K., Sundell, K.A., Hamidi, A., Skoog, I., Gustavsson, S. and Waern, M., 2018. Antidepressants and suicidal behaviour in late life: A prospective population-based study of use patterns in new users aged 75 and above. *European journal of clinical pharmacology*, 74(2), pp.201-208.
- Lawrence, B.J., Jayakody, D.M., Bennett, R.J., Eikelboom, R.H., Gasson, N. and Friedland, P.L., 2020. Hearing loss and depression in older adults: a systematic review and meta-analysis. *The Gerontologist*, 60(3), pp.e137-e154.
- Soysal, P., Veronese, N., Thompson, T., Kahl, K.G., Fernandes, B.S., Prina, A.M., Solmi, M., Schofield, P., Koyanagi, A., Tseng, P.T. and Lin, P.Y., 2017. Relationship between depression and frailty in older adults: A systematic review and meta-analysis. *Ageing research reviews*, 36, pp.78-87.

- Grief - when does it become pathological?
- Suicide in the elderly

MCQs

1. The features suggestive of depression-executive dysfunction syndrome would include all except:

- A. There is a long history of memory impairment and difficult with ADLs
- B. The patient complains of poor memory
- C. Assessment of cognitive function often results in 'don't know answers'
- D. The onset is fast
- E. There is often a history of depression or an identifiable precipitant

2. An 84 year old lady presents with severe depression. She had a myocardial infarction 3 months ago and her QTc is 490ms. Which antidepressant is the best choice?

- A. Sertraline
- B. Mirtazapine
- C. Paroxetine
- D. Citalopram
- E. Duloxetine

3. An 87 year old man has lost his wife recently. Which of the following clinical features would most suggest that this was an abnormal grief reaction?

- A. Loss of sleep
- B. Loss of appetite
- C. Laying the dining table for the deceased at meal times
- D. Anxiety
- E. Suicidal ideation

4. Which is not a feature of serotonin syndrome?

- A. Blurred vision
- B. Confusion
- C. Akathisia
- D. Elevated white cells
- E. Hypomimia

5. Which rating scale is most helpful in detecting depression in people with dementia?

- A. Cornell
- B. MMSE
- C. GDS
- D. AMTS
- E. Hamilton Rating Scale

6. You have a patient on lithium with a consistently elevated blood pressure. What is your most appropriate action?

- A. Start amiloride
- B. Lithium must be stopped
- C. Start furosemide
- D. Start lisinopril
- E. Start candesartan

[Additional Resources / Reading Materials](#)

Online:

- CPD Online: Quick bite: Suicide in the elderly, treating depression in later life, bereavement

Landmark studies

- Alexopoulos, G.S., Abrams, R.C., Young, R.C. and Shamoian, C.A., 1988. Cornell scale for depression in dementia. *Biological psychiatry*, 23(3), pp.271-284.
- Cipriani, A., Furukawa, T.A., Salanti, G., Geddes, J.R., Higgins, J.P., Churchill, R., Watanabe, N., Nakagawa, A., Omori, I.M., McGuire, H. and Tansella, M., 2009. Comparative efficacy and acceptability of 12 new-generation antidepressants: a multiple-treatments meta-analysis. *The lancet*, 373(9665), pp.746-758.
- Glassman, A.H., O'Connor, C.M., Califf, R.M., Swedberg, K., Schwartz, P., Bigger Jr, J.T., Krishnan, K.R.R., Van Zyl, L.T., Swenson, J.R., Finkel, M.S. and Landau, C., 2002. Sertraline treatment of major depression in patients with acute MI or unstable angina. *Jama*, 288(6), pp.701-709.
- Rush, A.J., Fava, M., Wisniewski, S.R., Lavori, P.W., Trivedi, M.H., Sackeim, H.A., Thase, M.E., Nierenberg, A.A., Quitkin, F.M., Kashner, T.M. and Kupfer, D.J., 2004. Sequenced treatment alternatives to relieve depression (STAR* D): rationale and design. *Controlled clinical trials*, 25(1), pp.119-142.
- Trivedi, M.H., Rush, A.J., Wisniewski, S.R., Nierenberg, A.A., Warden, D., Ritz, L., Norquist, G., Howland, R.H., Lebowitz, B., McGrath, P.J. and Shores-Wilson, K., 2006. Evaluation of outcomes with citalopram for depression using measurement-based care in STAR* D: implications for clinical practice. *American journal of Psychiatry*, 163(1), pp.28-40.

Journal Papers:

- Agüera-Ortiz, L., Claver-Martín, M.D., Franco-Fernández, M.D., López-Álvarez, J., Martín-Carrasco, M., Ramos-García, M.I. and Sánchez-Pérez, M., 2020. Depression in the elderly. Consensus statement of the Spanish Psychogeriatric Association. *Frontiers in Psychiatry*, 11.
- Aizenstein, H.J., Baskys, A., Boldrini, M., Butters, M.A., Diniz, B.S., Jaiswal, M.K., Jellinger, K.A., Kruglov, L.S., Meshandin, I.A., Mijajlovic, M.D. and Niklewski, G., 2016. Vascular depression consensus report—a critical update. *BMC medicine*, 14(1), pp.1-16.
- Andreas S, Dehoust M, Volkert J, Schulz H, Sehner S, Suling A, Wegscheider K, Ausín B, Canuto A, Crawford MJ, Da Ronch C. Affective disorders in the elderly in different European countries: Results from the MentDis_ICF65+ study. *PloS one*. 2019;14(11).
- Arthur, A., Savva, G.M., Barnes, L.E., Borjian-Borojony, A., Dening, T., Jagger, C., Matthews, F.E., Robinson, L. and Brayne, C., 2020. Changing prevalence and treatment of depression among older people over two decades. *The British Journal of Psychiatry*, 216(1), pp.49-54.
- Baez, S., Pinasco, C., Roca, M., Ferrari, J., Couto, B., García-Cordero, I., Ibañez, A., Cruz, F., Reyes, P., Matallana, D. and Manes, F., 2019. Brain structural correlates of executive and social cognition profiles in behavioral variant frontotemporal dementia and elderly bipolar disorder. *Neuropsychologia*, 126, pp.159-169.
- Büchtemann, D., Luppá, M., Bramesfeld, A. and Riedel-Heller, S., 2012. Incidence of late-life depression: a systematic review. *Journal of affective disorders*, 142(1-3), pp.172-179.
- Cai W, Mueller C, Shetty H, Perera G, Stewart R. Predictors of mortality in people with late-life depression: A retrospective cohort study. *Journal of Affective Disorders*. 2020 Jan 10.
- Caixeta, L., Soares, V.L., Vieira, R.T., Soares, C.D., Caixeta, V., Ferreira, S.B. and Aversi-Ferreira, T.A., 2017. Executive function is selectively impaired in old age bipolar depression. *Frontiers in Psychology*, 8, p.194.
- Cattell, H. 2000. Suicide in the elderly. *Advances in Psychiatric Treatment*, 6(2), 102-108.
- Cipriani, A., Furukawa, T.A., Salanti, G., Chaimani, A., Atkinson, L.Z., Ogawa, Y., Leucht, S., Ruhe, H.G., Turner, E.H., Higgins, J.P. and Egger, M., 2018. Comparative efficacy and acceptability of 21 antidepressant drugs for the acute treatment of adults with major depressive disorder: a systematic review and network meta-analysis. *Focus*, 16(4), pp.420-429.
- Conwell, Y., 2014. Suicide later in life: challenges and priorities for prevention. *American Journal of Preventive Medicine*, 47(3), pp.S244-S250.

- Cooper, C., Katona, C., Lyketsos, K., Blazer, D., Brodaty, H., Rabins, P., de Mendonça Lima, C.A. and Livingston, G., 2011. A systematic review of treatments for refractory depression in older people. *American Journal of psychiatry*, 168(7), pp.681-688.
- Djernes, J.K., 2006. Prevalence and predictors of depression in populations of elderly: a review. *Acta Psychiatrica Scandinavica*, 113(5), pp.372-387.
- Dols, A., Kessing, L.V., Strejilevich, S.A., Rej, S., Tsai, S.Y., Gildengers, A.G., Almeida, O.P., Shulman, K.I., Sajatovic, M. and International Society for Bipolar Disorders Task Force for Older Adults with Bipolar Disorder, 2016. Do current national and international guidelines have specific recommendations for older adults with bipolar disorder? A brief report. *International journal of geriatric psychiatry*, 31(12), pp.1295-1300.
- Draper, B. M. 2014. Suicidal behaviour and suicide prevention in later life. *Maturitas*, 79(2), 179-183.
- Fancourt, D. and Tymoszuk, U., 2019. Cultural engagement and incident depression in older adults: evidence from the English Longitudinal Study of Ageing. *The British Journal of Psychiatry*, 214(4), pp.225-229.
- Gadzhanova, S., Roughead, E.E. and Pont, L.G., 2018. Antidepressant switching patterns in the elderly. *International psychogeriatrics*, 30(9), pp.1365-1374.
- Geduldig, E.T. and Kellner, C.H., 2016. Electroconvulsive therapy in the elderly: new findings in geriatric depression. *Current psychiatry reports*, 18(4), p.40.
- Haigh, E.A., Bogucki, O.E., Sigmon, S.T. and Blazer, D.G., 2018. Depression among older adults: a 20-year update on five common myths and misconceptions. *The American Journal of Geriatric Psychiatry*, 26(1), pp.107-122.
- Hegeman, J.M., Kok, R.M., Van der Mast, R.C. and Giltay, E.J., 2012. Phenomenology of depression in older compared with younger adults: meta-analysis. *The British Journal of Psychiatry*, 200(4), pp.275-281.
- John A, Patel U, Rusted J, Richards M, Gaysina D. Affective problems and decline in cognitive state in older adults: a systematic review and meta-analysis. *Psychological medicine*. 2019 Feb;49(3):353-65.
- Kok, R.M. and Reynolds, C.F., 2017. Management of depression in older adults: a review. *Jama*, 317(20), pp.2114-2122.
- Lawrence, B.J., Jayakody, D.M., Bennett, R.J., Eikelboom, R.H., Gasson, N. and Friedland, P.L., 2020. Hearing loss and depression in older adults: a systematic review and meta-analysis. *The Gerontologist*, 60(3), pp.e137-e154.

- Lisanby, S.H., McClintock, S.M., Alexopoulos, G., Bailine, S.H., Bernhardt, E., Briggs, M.C., Cullum, C.M., Deng, Z.D., Dooley, M., Geduldig, E.T. and Greenberg, R.M., 2020. Neurocognitive effects of combined electroconvulsive therapy (ECT) and venlafaxine in geriatric depression: phase 1 of the PRIDE study. *The American Journal of Geriatric Psychiatry*, 28(3), pp.304-316.
- Liu, L., Gou, Z. and Zuo, J., 2016. Social support mediates loneliness and depression in elderly people. *Journal of health psychology*, 21(5), pp.750-758.
- Luppá, M., Sikorski, C., Luck, T., Ehreke, L., Konnopka, A., Wiese, B., Weyerer, S., König, H.H. and Riedel-Heller, S.G., 2012. Age-and gender-specific prevalence of depression in latest-life—systematic review and meta-analysis. *Journal of affective disorders*, 136(3), pp.212-221.
- Mallery L, MacLeod T, Allen M, McLean-Veysey P, Rodney-Cail N, Bezanson E, Steeves B, LeBlanc C, Moorhouse P. Systematic review and meta-analysis of second-generation antidepressants for the treatment of older adults with depression: questionable benefit and considerations for frailty. *BMC geriatrics*. 2019 Dec;19(1):1-1.
- Mann, E., Hinrichsen, G.A. and Baharlou, S., 2020. Depression in Older Adults: Principles of Diagnosis and Management. In *Geriatric Practice* (pp. 213-221). Springer, Cham.
- McDonald, W.M., Hermida, A., Petrides, G. and Kellner, C., 2017. Update on New Research and the Clinical Practice of ECT in the Elderly. *The American Journal of Geriatric Psychiatry*, 25(3), p.S25.
- McDOUGALL, F.A., Kvaal, K., Matthews, F.E., Paykel, E., Jones, P.B., Dewey, M.E. and Brayne, C., 2007. Prevalence of depression in older people in England and Wales: the MRC CFA Study. *Psychological medicine*, 37(12), p.1787.
- Morin RT, Nelson C, Bickford D, Insel PS, Mackin RS. Somatic and anxiety symptoms of depression are associated with disability in late life depression. *Aging & mental health*. 2019 Apr 3:1-4.
- Morse, J.Q. and Robins, C.J., 2005. Personality–life event congruence effects in late-life depression. *Journal of affective disorders*, 84(1), pp.25-31.
- Nair P, Bhanu C, Frost R, Buszewicz M, Walters KR. A Systematic Review of Older Adults' Attitudes towards Depression and its Treatment. *The Gerontologist*. 2020 Jan 24;60(1):e93-104.
- Naismith, S.L., Norrie, L.M., Mowszowski, L. and Hickie, I.B., 2012. The neurobiology of depression in later-life: clinical, neuropsychological, neuroimaging and pathophysiological features. *Progress in neurobiology*, 98(1), pp.99-143.
- Obbels J, Vansteelandt K, Verwijk E, Dols A, Bouckaert F, Oudega ML, Vandenbulcke M, Stek M, Sienaert P. MMSE changes during and after ECT in late-life depression: a prospective study. *The American Journal of Geriatric Psychiatry*. 2019 Sep 1;27(9):934-44.

- Prakash, A., 2019. Psychology of Depression in Elderly: A Review. *EC Psychology and Psychiatry*, 8, pp.263-272.
- Pruckner, N. and Holthoff-Detto, V., 2017. Antidepressant pharmacotherapy in old-age depression—a review and clinical approach. *European Journal of Clinical Pharmacology*, 73(6), pp.661-667.
- Reppermund, S., 2016. Depression in old age—the first step to dementia?. *The Lancet Psychiatry*, 3(7), pp.593-595.
- Richards, F., & Curtice, M. 2011. Mania in late life. *Advances in Psychiatric Treatment*, 17(5), 357-364.
- Rodda, J., Walker, Z., & Carter, J. 2011. Depression in older adults. *BMJ*, 343.
- Rossom RC, Simon GE, Coleman KJ, Beck A, Oliver M, Stewart C, Ahmedani B. Are wishes for death or suicidal ideation symptoms of depression in older adults?. *Aging & mental health*. 2019 Jul 3;23(7):912-8.
- Rudd KB, Breen R, Srinivasan S, Hrisko S. Suicide in late life: Collaborative approaches for assessment, prevention and treatment: Session 202. *The American Journal of Geriatric Psychiatry*. 2019 Mar 1;27(3):S13-4.
- Shulman, K.I., Almeida, O.P., Herrmann, N., Schaffer, A., Strejilevich, S.A., Paternoster, C., Amodeo, S., Dols, A. and Sajatovic, M., 2019. Delphi survey of maintenance lithium treatment in older adults with bipolar disorder: an ISBD task force report. *Bipolar disorders*, 21(2), pp.117-123.
- Sobieraj DM, Martinez BK, Hernandez AV, Coleman CI, Ross JS, Berg KM, Steffens DC, Baker WL. Adverse effects of pharmacologic treatments of major depression in older adults. *Journal of the American Geriatrics Society*. 2019 Aug;67(8):1571-81.
- Soysal, P., Veronese, N., Thompson, T., Kahl, K.G., Fernandes, B.S., Prina, A.M., Solmi, M., Schofield, P., Koyanagi, A., Tseng, P.T. and Lin, P.Y., 2017. Relationship between depression and frailty in older adults: A systematic review and meta-analysis. *Ageing research reviews*, 36, pp.78-87.
- Van Assche, L., Van de Ven, L., Vandenbulcke, M. and Luyten, P., 2019. Ghosts from the past? The association between childhood interpersonal trauma, attachment and anxiety and depression in late life. *Aging & Mental Health*, pp.1-8.
- Volkert J, Härter M, Dehoust MC, Ausín B, Canuto A, Da Ronch C, Suling A, Grassi L, Munoz M, Santos-Olmo AB, Sehner S. The role of meaning in life in community-dwelling older adults with depression and relationship to other risk factors. *Aging & mental health*. 2019 Jan 2;23(1):100-6.

- Wei J, Hou R, Zhang X, Xu H, Xie L, Chandrasekar EK, Ying M, Goodman M. The association of late-life depression with all-cause and cardiovascular mortality among community-dwelling older adults: systematic review and meta-analysis. *The British Journal of Psychiatry*. 2019 Aug;215(2):449-55.

Books:

- Denning T., Thomas A., 2013. *The Oxford Textbook of Old Age Psychiatry*, 2nd edition. Oxford University Press.
- Stahl, SM, 2017. *Prescriber's Guide: Stahl's Essential Psychopharmacology*, 6th edition Cambridge University Press.
- Taylor, D., Barnes, T., Young, A., 2018. *The Maudsley Prescribing Guidelines in Psychiatry*, 13th edition. Blackwell-Wiley.(sections on mood disorders including prescribing in older adults).
- World Health Organisation, 1992. *ICD-10: The ICD-10 Classification of Mental and Behavioural Disorders : Clinical Descriptions and Diagnostic Guidelines*. WHO.

Session 6: Psychosis in the Older Person

Learning Objectives

- The overall aim of the sessions is for the trainees to gain an overview of psychosis in later life.
- By the end of the session trainees should:
 - Understand the epidemiology of psychosis and psychotic disorders in the older person.
 - Understand the aetiology of psychosis in the older person.
 - Understand how psychosis presents in the older person, the classification of disorders, the basic assessment process and the principles of treatment of psychosis and psychotic disorders

Curriculum Links

- Old Age Section of the MRCPsych Curriculum: 8.3, 8.4, 8.5, 8.6, 8.7, 8.8, 8.9

Expert Led Session

- A Consultant led session based on the learning objectives listed above.

Case Presentation

- A case involving an older person presenting with probable psychosis. Please consider the learning objectives above.

Journal Club Presentation

- Howard, R., Cort, E., Bradley, R., Kelly, L., Bentham, P., Ritchie, C., Reeves, S., Fawzi, W., Livingston, G., Sommerlad, A. and Oomman, S., 2018. Antipsychotic treatment of very late-onset schizophrenia-like psychosis: a randomised controlled double-blind trial. *The Lancet Psychiatry*.
- Stafford, J., Dykxhoorn, J., Sommerlad, A., Dalman, C., Kirkbride, J. and Howard, R., 2020. Association between risk of dementia and very late-onset schizophrenia-like psychosis: a Swedish population-based cohort study.
- Zhang, H., Wang, L., Fan, Y., Yang, L., Wen, X., Liu, Y. and Liu, Z., 2019. Atypical antipsychotics for Parkinson's disease psychosis: a systematic review and meta-analysis. *Neuropsychiatric disease and treatment*, 15, p.2137.

'555' Topic (5 slides with no more than 5 bullet points per slide)

- Antipsychotics and the elderly – factors to consider when prescribing
- Schizophrenia in adults vs older adults - the key differences.
- Charles Bonnet syndrome

1. A 76 year old lady is diagnosed with 'late paraphrenia'. Which of the following delusions is the GP most likely to find?

- A. Hypochondriachal
- B. Delusions of misidentification
- C. Religious delusions
- D. Delusions of reference
- E. Persecutory delusions

2. Very late onset schizophrenia is characterised by onset after:

- A. 40 years
- B. 60 years
- C. 65 years
- D. 70 years
- E. 80 years

3. Which antipsychotic is most likely to cause postural hypotension:

- A. Aripiprazole
- B. Risperidone
- C. Haloperidol
- D. Quetiapine
- E. Sulpiride

4. Which of the following drugs should not be used in renal failure?

- A. Amisulpride
- B. Aripiprazole
- C. Chlorpromazine
- D. Olanzapine
- E. Quetiapine

5. 'Sensitivity to antipsychotics' is a core feature of which disorder?

- A. Alzheimer's Disease
- B. Dementia with Lewy Bodies
- C. Late onset Schizophrenia
- D. Organic mood disorder
- E. Huntington's Disease

Online:

- RCPsych CPD online. The management of hyperprolactinemia in psychiatric practice, psychotropic medication and the heart

Landmark papers

- Howard, R., Cort, E., Bradley, R., Harper, E., Kelly, L., Bentham, P., Ritchie, C., Reeves, S., Fawzi, W., Livingston, G. and Sommerlad, A., 2018. Antipsychotic treatment of very late-onset schizophrenia-like psychosis (ATLAS): a randomised, controlled, double-blind trial. *The Lancet Psychiatry*, 5(7), pp.553-563.
- Howard, R., Rabins, P. V., Seeman, M. V., & Jeste, D. V. 2000. Late-onset schizophrenia and very-late-onset schizophrenia-like psychosis: an international consensus. *American Journal of Psychiatry*.

Journal Papers:

- Almeida, O.P., Ford, A.H., Hankey, G.J., Yeap, B.B., Golledge, J. and Flicker, L., 2019. Risk of dementia associated with psychotic disorders in later life: the health in men study (HIMS). *Psychological medicine*, 49(2), pp.232-242.
- Andreas, S., Schulz, H., Volkert, J., Dehoust, M., Sehner, S., Suling, A., Ausín, B., Canuto, A., Crawford, M., Da Ronch, C. and Grassi, L., 2017. Prevalence of mental disorders in elderly people: the European MentDis_ICF65+ study. *The British Journal of Psychiatry*, 210(2), pp.125-131.
- Bartels, S.J., Fortuna, K.L. and Naslund, J.A., 2018. Serious Mental Disorders in Older Adults: Schizophrenia and Other Late-Life Psychoses. *Aging and Mental Health*, pp.241-280.
- Ferenczi, E.A., Erkinen, M.G., Feany, M.B., Fogel, B.S. and Daffner, K.R., 2020. New-Onset Delusions Herald an Underlying Neurodegenerative Condition: A Case Report and Review of the Literature. *The Journal of Clinical Psychiatry*, 81(2).
- Fischer, C.E., Ismail, Z., Youakim, J.M., Creese, B., Kumar, S., Nuñez, N., Darby, R.R., Di Vita, A., D'Antonio, F., de Lena, C. and McGeown, W.J., 2019. Revisiting Criteria for Psychosis in Alzheimer's Disease and Related Dementias: Toward Better Phenotypic Classification and Biomarker Research. *Journal of Alzheimer's Disease*, (Preprint), pp.1-14.
- Lange, S.M., Meesters, P.D., Stek, M.L., Wunderink, L., Penninx, B.W. and Rhebergen, D., 2019. Course and predictors of symptomatic remission in late-life schizophrenia: A 5-year follow-up study in a Dutch psychiatric catchment area. *Schizophrenia research*, 209, pp.179-184.
- Lapid, M.I. and Ho, J.B., 2020. Challenging our beliefs about delusional disorder in late life. *International Psychogeriatrics*, 32(4), pp.423-425.

- Louhija, U.M., Saarela, T., Juva, K. and Appelberg, B., 2017. Brain atrophy is a frequent finding in elderly patients with first episode psychosis. *International psychogeriatrics*, 29(11), pp.1925-1929.
- Maglione, J.E., Thomas, S.E. and Jeste, D.V., 2014. Late-onset schizophrenia: do recent studies support categorizing LOS as a subtype of schizophrenia?. *Current opinion in psychiatry*, 27(3), p.173.
- Rossi, M., Farcy, N., Starkstein, S.E. and Merello, M., 2020. Nosology and Phenomenology of Psychosis in Movement Disorders. *Movement disorders clinical practice*.
- Rothenberg, K.G. and Rajaram, R., 2019. Advances in Management of Psychosis in Neurodegenerative Diseases. *Current treatment options in neurology*, 21(1), p.3.
- Suen, Y.N., Wong, S.M., Hui, C.L., Chan, S.K., Lee, E.H., Chang, W.C. and Chen, E.Y., 2019. Late-onset psychosis and very-late-onset-schizophrenia-like-psychosis: an updated systematic review. *International Review of Psychiatry*, 31(5-6), pp.523-542.
- Swann P, O'Brien JT. Management of visual hallucinations in dementia and Parkinson's disease. *International psychogeriatrics*. 2019 Jun;31(6):815-36.
- Tampi, R.R., Young, J., Hoq, R., Resnick, K. and Tampi, D.J., 2019. Psychotic disorders in late life: a narrative review. *Therapeutic advances in psychopharmacology*, 9, p.2045125319882798.
- Van Assche, L., Morrens, M., Luyten, P., Van de Ven, L. and Vandenbulcke, M., 2017. The neuropsychology and neurobiology of late-onset schizophrenia and very-late-onset schizophrenia-like psychosis: a critical review. *Neuroscience & Biobehavioral Reviews*, 83, pp.604-621.
- Van Assche, L., Van Aubele, E., Van de Ven, L., Bouckaert, F., Luyten, P. and Vandenbulcke, M., 2019. The neuropsychological profile and phenomenology of late onset psychosis: a cross-sectional study on the differential diagnosis of very-late-onset schizophrenia-like psychosis, dementia with Lewy bodies and Alzheimer's type dementia with psychosis. *Archives of Clinical Neuropsychology*, 34(2), pp.183-199.
- Zharkova, T. and Kyomen, H.H., 2018. Treatment Dilemmas: Managing Antipsychotic Medication Risks in Elderly with Major Neurocognitive Disorder, Stroke and Psychosis. *The American Journal of Geriatric Psychiatry*, 26(3), pp.S100-S101.

Guidelines:

- Psychosis and schizophrenia in adults: prevention and management. NICE guidelines [CG178]

Books:

- Denning T., Thomas A., 2013. *The Oxford Textbook of Old Age Psychiatry*, 2nd edition. Oxford University Press.

- Vannorsdall, T.D. and Schretlen, D.J., 2019. Late-onset schizophrenia. In Handbook on the neuropsychology of Aging and Dementia (pp. 711-725). Springer, Cham.
- Stahl, SM, 2017. Prescriber's Guide: Stahl's Essential Psychopharmacology, 6th edition Cambridge University Press.
- Taylor, D., Barnes, T., Young, A., 2018. The Maudsley Prescribing Guidelines in Psychiatry, 13th edition. Blackwell-Wiley.
- World Health Organisation, 1992. ICD-10: The ICD-10 Classification of Mental and Behavioural Disorders: Clinical Descriptions and Diagnostic Guidelines. WHO.

MCQ answers

Mood disorders

1. **A** - a long history of memory impairment and difficulty with ADLS more suggestive of dementia
2. **A** – sertraline as supported by the SADHART trial results
3. **E**
4. **E**
5. **A**
6. **A**

Psychosis

1. **E**
2. **B**
3. **D**
4. **A**
5. **B**

Across the Ages

To follow

Forensic

Session 3: Too mad to murder?

Learning Objectives

- To develop an understanding of the role of mental disorder in offending
- To develop an understanding of the frequency of and types of offences committed by those with serious mental illness
- To understand the role of special syndromes in offences
- To develop an understanding of vulnerability and suggestibility in mentally disordered offenders

Curriculum Links

12.1 Relationship between crime and mental disorder

12.1.2 The relationship between specific mental disorders and crime: substance misuse; epilepsy; schizophrenia; bipolar affective disorder; neuro-developmental disorders; personality disorders

12.1.3 Special syndromes: morbid jealousy, erotomania, Munchausen and Munchausen by proxy

12.1.5 Effect of victimisation and vulnerability: anxiety states including post-traumatic stress disorder; suggestibility; anger and aggressive behaviour. Effect of compensation on presentation

Expert Led Session

'Too mad to murder?' to include:

- Substance Misuse
- Epilepsy

- Schizophrenia
- Bipolar affective disorder
- Neuro-developmental disorders
- Personality disorders

Case Presentation

Case presentation on 'a special syndrome in relation to forensic psychiatry'. To include either morbid jealousy, erotomania, Munchausen or Munchausen by proxy.

Options for case presentation:

- If trainee has a suitable case of a special syndrome then they may present this.
- The trainee can come to the Edenfield Centre where a suitable case can be found for them – to access case notes and / or meet patient (if appropriate)
- To use 'The Allitt inquiry' as the basis of the case presentation. (Munchausen by proxy)

Journal Club Presentation

Key points to be summarised from the following three papers:

- Rose J, Cutler C, Tresize K et al (2008) Individuals with an intellectual disability who offend, British Journal of Developmental Disabilities 106, 19 – 30
http://www.google.co.uk/url?sa=t&rct=j&q=&esrc=s&frm=1&source=web&cd=1&cad=rja&uact=8&sqi=2&ved=0CCUQFjAA&url=http%3A%2F%2Fwww.researchgate.net%2Fpublication%2F228505583_Individuals_with_an_intellectual_disability_who_offend%2Flinks%2F0deec51817f57baef7000000&ei=3YngU_Pil-nb7Aazh4DABq&usq=AFQjCNEg9xYeimpqgJchT70fngkh2vkPTA&sig2=KXDBJ1CC_DT2OPQG6mr2KA
- Fazel S, Wolf A, Chang Z et al (2015). Depression and violence: a Swedish population study. Lancet Psychiatry 2: 224 – 32
- Elbogen EB & Johnson SC (2009) The intricate link between violence and mental disorder: results from the national epidemiological survey on alcohol and related conditions. Archives of General Psychiatry 66(2): 152 – 161

<http://www.ncbi.nlm.nih.gov/pubmed/19188537>

'555' Topic (5 slides with no more than 5 bullet points per slide)

- Human rights legislation – articles 5 / 6 / 8
- Ethics

MCQs

1. Which is the biggest risk factor for violence in psychosis?
 - A. Non-compliance with medication
 - B. Co-morbid personality disorder
 - C. Homelessness
 - D. Unemployment
 - E. Co-morbid substance misuse
2. With respect to Munchausen's by Proxy, which of the following is incorrect?
 - A. More common in mothers
 - B. The annual incidence of fabricated or induced illness in children under 16 is 0.5 per 100,000
 - C. There is no clear relationship with any specific mental disorder
 - D. 50% perpetrators had a personality disorder
 - E. 21% have a history of alcohol and / or drug misuse
3. Which of the following regarding mood disorder and violence is incorrect?
 - A. The prevalence of depression in male prisoners is 10%
 - B. The prevalence of depression in female prisoners is 25%
 - C. Manic patients are likely to show aggression and violence associated with admission to hospital
 - D. 7% homicide perpetrators have a lifetime diagnosis of mood disorder
 - E. Most perpetrators of homicide-suicide are male
4. Which is the correct statement relating to substance use and the MacArthur Violence Study?
 - A. Substance use increases the rate of violence among both those with and without mental illness
 - B. The rate of violence for those with a mental disorder and no substance use is 25%
 - C. The rate of violence for those with a mental disorder and substance use is 50%
 - D. Substance use is a protective factor for violence

E. The highest rate of violence was for those with mood disorder and substance use

5. Which is the incorrect statement about epilepsy and offending?

A. Ictal violence is more likely in complex partial seizures

B. Most offending occurs in post-ictal or inter-ictal period

C. Violence in epilepsy is usually a feature of the disease

D. The prevalence of epilepsy in prisoners is 1 – 2%

E. The prevalence of epilepsy in the general population is 0.5 – 1%

EMI Questions

Fire Setting:

A. Crime concealment

B. Financial compensation

C. Suicidal

D. Extremism

E. Vandalism

F. Psychosis

H. Pyromania

Match the most-likely motivation for fire-setting with the clinical scenario below.

1. Wayne is a 14 year old who whilst truanting from school with a gang of boys sets fire to an abandoned warehouse. He waits around for the fire service to arrive and watches from a safe distance as they put the fire out.

2. Vincent is a 48 year old man with Asperger's Disorder. He has a history of setting fires when he is angry. He enjoys looking at how things burn. He is upset by another resident shouting at him and so set a fire. He feels an inner tension that is relieved when he has set the fire. He calls the fire brigade and becomes excited when they arrive.

3. Stephanie sets fire to a university research laboratory, where she believes the researchers are carrying out experiments on elephants. Two weeks ago she suddenly realised that the

University were dissecting elephant trunks in order to test the effects of snorting cocaine so that the Government could develop a synthetic drug to distribute in the community.

4. Alison is a 50 year old woman who has recently separated from her husband after he left her for another woman. Divorce proceedings have begun and she is concerned that she may have to leave the family home because she can't afford to pay the mortgage. She is depressed with low mood, poor sleep, anhedonia and poor concentration. She feels that if she loses her home she won't have anything to live for. She sets fire to her house using petrol in 3 seats in the living room, hallway and upstairs bedroom. She calls the fire brigade from her mobile phone in the garden.

Human Rights:

- A. Article 2
- B. Article 3
- C. Article 5
- D. Article 6
- E. Article 8
- F. Article 9
- G. Article 12**

These Articles of the European Convention of Human Rights (ECHR) are important in the detention of mentally-disordered offenders. Match the correct Article with the freedom or right it describes.

1. Right to respect for private and family life
2. Prohibition of torture
3. Right to marry
4. Right to life
5. Right to liberty and security
6. Freedom of thought, conscience and religion
7. Right to a fair trial

[Additional Resources / Reading Materials](#)

Books

- Chapters 14, 16, 17, 18, & 26 in 'Forensic Psychiatry: Clinical and ethical issues' Gunn J & Taylor P, (2013) CRC Press
- Chapters 7, 8 & 9 in 'Practical Forensic Psychiatry,' Clark T & Rooprai DS (2011) Hodder Arnold
- Chapter 3 in 'Oxford Specialist Handbook: Forensic Psychiatry,' Eastman N, Adshead G, Fox S et al (2012) Oxford Medical Publishing

E-Learning

- RCPsych CPD online: 'FREDA – a human rights-based approach to clinical practice'
- RCPsych CPD online: 'Morbid jealousy'
- RCPsych CPD online: 'Understanding and safely managing paranoid personality disorder'

Journal Articles

- Arsenault L, Moffit T, Caspi A et al (2000) Mental disorders and violence: results from the Dunedin study. Archives of General Psychiatry 57: 979 – 986
- Barrowcliff AL & Haddock G (2006) The relationship between command hallucinations and factors of compliance: a critical review of the literature. Journal of forensic psychiatry and psychology 17(2): 266 – 298
- Booles CN, Neale BA, Meadow SR (1994) Munchausen syndrome by proxy: a study of psychopathology. Child abuse and neglect 18: 773 – 788
- Fazel S, Langstrom N, Hjern A et al (2009) Schizophrenia, substance abuse, and violent crime. Journal of the American Medical Association 301(19): 2016 – 2023
- Gudjonsson GH & Henry L. (2003) Child and adult witnesses with intellectual disability: the importance of suggestibility Legal and Criminological Psychology 8(2), 241 – 252
- Large M, Smith G, Swinson N et al (2008) Homicide due to mental disorder in England and Wales over 50 years. British Journal of Psychiatry 193: 130 – 133
- Newhill CE, Eack SM & Mulvey EP (2009) Violent behavior in borderline personality disorder. Journal of Personality Disorders 23: 541 – 554

- Nielson O & Large M (2010) Rates of homicide during the first episode of psychosis and after treatment: a systematic review and meta-analysis *Schizophrenia Bulletin* 36(4): 702 – 712
- Roberts ADL & Coid JW (2010) Personality disorder and offending behaviour: findings from the national survey of male prisoners in England and Wales. *Journal of forensic psychiatry and psychology* 21: 221 – 237
- Shaw J, Amos T, Hunt IM et al (2004) Mental illness in people who kill strangers: longitudinal study and national clinical survey. *British Medical Journal* 328: 734 – 737
- Shaw J, Amos T, Hunt IM et al (2006) Rates of mental disorder in people convicted of homicide. *British Journal of Psychiatry* 188: 143 - 147
- Swanson JW, Holzer CE, Ganju VK, Jono R (1990) Violence and psychiatric disorder in the community: evidence from the epidemiological catchment area survey *Hospital and Community Psychiatry* 41, 761 – 70
- Tihonen J, Isohanni M, Rasanen P et al (1997) Specific major mental disorders and criminality: a 26 year prospective study of the 1966 northern Finland birth cohort. *American Journal of Psychiatry* 154: 840 – 845

ID
Session 3: Behavioural Issues in Intellectual Disability
Learning Objectives
<ul style="list-style-type: none"> • Understanding challenging behaviour and awareness of methods of recording/ assessing • Aetiology of challenging behaviours • Management options
Curriculum Links
<p>13.1 Services</p> <p>13.1.2 The provision of specialist psychiatric services for people with intellectual disability</p> <p>13.2.1 The factors which might account to the observed high rates of psychiatric behavioural disorders in this group</p> <p>13.3.2 The presentation and diagnosis of psychiatric illness and behavioural disorder in people with intellectual disability, including the concept of diagnostic overshadowing</p>
Expert Led Session
<ul style="list-style-type: none"> • Challenging Behaviour Talk
Case Presentation
<ul style="list-style-type: none"> • Case presentation of local patient with intellectual disability presenting with behavioural problems, identified by tutor or specialist in post (this does not have to be an inpatient and discussion with the local ID team may be appropriate in advance to identify such a case). Brief discussion on aetiology as applicable to the case in a formulation type summary
Journal Club Presentation
<p>Please select one of the following papers:</p> <ul style="list-style-type: none"> • Unwin G.L. and Deb S. (2008) A multi-centre audit of the use of medication for the management of behavioural problems in adults with intellectual disabilities. British Journal of Learning Disabilities, 36, 2, 140-143 • Cooper S.A. Melville C.A. and Enfield S.L. (2003) Psychiatric diagnosis, intellectual disabilities and Diagnostic Criteria for Psychiatric Disorders for Use with Adults with Learning Disabilities/Mental Retardation (DC-LD). Journal of Intellectual Disability Research 47, supplement one, 3-15. • Group-based cognitive-behavioural anger management for people with mild to moderate intellectual disabilities: cluster randomised controlled trial BJP October 2013 203:288-296;

'555' Topics (5 slides on each topic with no more than 5 bullet points)

Please select one of the following:

- Review of Frith Guidelines on management of Patients with ID that present with Aggressive or Self Injurious behaviours. (Read the Guidelines in particular the flow charts)
- Describe challenging behaviour and the various phases of the cycle of challenging behaviour (Focus on nature of behaviours, communication ability of the patient, issues of any change).
- Formal Assessment of a behavioural problem with a view to intervention. (You can discuss this with your local ID team to guide with the task). Steps involved, would include ABC charts or functional assessments and basic behavioural interventions

MCQ's

1. Causes of challenging behaviour in a person with learning disability:

- A. Pain
- B. Overstimulation
- C. Under stimulation
- D. Wanting attention
- E. All of the above

2. The following statements are true of factors increasing challenging behaviours in a person with learning disability except which option?

- A. Undetected physical illness
- B. Communication problems
- C. Underlying mental illness
- D. Environmental issues
- E. Problem solving ability

3. Inappropriate behaviours may be maintained by re-enforcement from others. Which of the following is a process that helps to identify factors maintaining that behaviour?

- A. Functional analysis
- B. Statistical analysis
- C. Procedural analysis**
- D. Behavioural analysis

EMI Questions

Match each of the following psychological strategies to their possible effects:

A. Proactive Strategies

B. Positive Programming

C. Focused Support

D. Reactive Strategies

1. Systematic instructions given for greater skills and competence development which improves social integration
2. To produce rapid results and reduce reactive strategies
3. Designed to manage the behaviours at the time they occur
4. To produce change over time

Additional Resources/ Reading Materials

E-Learning

www.LD-Medication.bham.ac.uk

British Psychological Society and Royal College of Psychiatrists (BPS & RCPsych, 2006).
Challenging behaviour: a unified approach. Available:

<http://www.rcpsych.ac.uk/pdf/23%2009%202011%20LD%20PSYCH%20READING%20LIST.pdf>

Psychotherapy
Session 3: Psychological approaches to Depression
Learning Objectives
<p>To increase awareness of the psychological aspects of Depressive Disorder.</p> <p>To have an introductory knowledge of the main psychological models for depression.</p> <p>To have an overview of psychological treatments for Depression</p>
Curriculum Links
1.1, 1.2, 1.3, 1.3.4, 2.3, 2.4, 2.6, 2.8, 6.1, 7.1.1, 9, 14
Expert Led Session
An overview of psychological therapies for Depressive Disorder
Case Presentation
<p>This should be of a patient with depression, not necessarily one who is in / has had therapy.</p> <p>There should be sufficient background history to generate a discussion about the psycho-social factors in aetiology</p>
Journal Club Presentation
<p>The paper should preferably be selected in discussion with the chair / presenter of the expert led session</p> <ul style="list-style-type: none"> • Driessen et al (2015) The efficacy of short-term psychodynamic psychotherapy for depression: A meta-analysis update Clinical Psychology Review 42: 1-15 • Gottems Bastos et al (2015) The efficacy of long-term psychodynamic psychotherapy, fluoxetine and their combination in the outpatient treatment of depression Psychotherapy Research 25(5): 612-624 • (Other paper suggested by expert if applicable – contact the person doing ELS)
'555' Topics (5 slides on each topic with no more than 5 bullet points)
<p>Select one of the following:</p> <ul style="list-style-type: none"> • Psychological factors in the aetiology of depression

- Psychological symptoms of depression
- Current psychological treatments for depression recommended by NICE

MCQs

1. NICE guidance (CG90):
 - A. Recommends Computerised CBT for mild-moderate depression
 - B. Recommends Psychotherapy for severe depression
 - C. Advises not combining medication with psychological therapies
 - D. Recommends Cognitive therapy for relapse prevention
 - E. Defines Short-term Psychodynamic Psychotherapy as 10-15 sessions over 3-4 months
2. Cognitive Therapy:
 - A. Is originally based on the work of Judith Beck
 - B. Identifies Cognitive Errors that lead to or maintain depressive thoughts
 - C. Focuses on non-conscious thought content
 - D. Is enhanced by concurrent antidepressant treatment
 - E. Should not be used in older patients
3. Psychodynamic Therapies:
 - A. Have no evidence base for effectiveness
 - B. Are based on the model of the mind put forward by Freud
 - C. Seek to eradicate a patient's defences
 - D. Were among the first to link depression to loss
 - E. Focus on the past
4. Psychological factors in the aetiology of depression include
 - A. Parental indifference
 - B. Social circumstance
 - C. Maternal Depression
 - D. Cognitive biases or distortions
 - E. Bereavement
5. Evidence of effectiveness in the treatment of depression exists for:

- A. Psychoanalytic therapy
- B. Interpersonal Therapy
- C. 'Low intensity' therapy in IAPT
- D. Mentalization based CBT
- E. EMDR

Additional Resources / Reading Materials

Sigmund Freud "Mourning and Melancholia" (1917 [1915]) Standard Edition **14**: 237-258

Aaron Beck "Cognitive Therapy and the Emotional Disorders" 1976

Substance Misuse

Session 3: Diagnosis and Management of People with co-occurring mental health and alcohol/drug use conditions

Learning Objectives

- To develop understanding of key aspects in the diagnosis and treatment of patients with co-occurring mental health and alcohol/drug use conditions
- To increase awareness of complications with pharmacological treatment in patients with co-occurring mental health and alcohol/drug use conditions
- To develop knowledge of risk issues in people with co-occurring mental health and alcohol/drug use conditions
- To understand how local services are implemented to manage people with co-occurring mental health and alcohol/drug use conditions

Curriculum Links

- 11.1 Basic pharmacology and epidemiology
- 11.5 Effect of drug and alcohol use on psychiatric illness

Expert Led Session

- Diagnosis and Management of People with co-occurring mental health and alcohol/drug use conditions
 - Concepts
 - Epidemiology
 - Case studies
- There is sufficient material for a separate lecture if spare capacity on the programme.
- If feasible joint presentation by Substance Misuse Consultant and General Adult Psychiatrist for both sessions

Case Presentation

- Examine risk aspects of people with co-occurring mental health and alcohol/drug use conditions
- Relationship of the substance use to development of the symptoms

Journal Club Presentation

- Asher CJ, Gask L. (2010) Reasons for illicit drug use in people with schizophrenia: Qualitative study. *BMC Psychiatry*, 10:94.
- Chitty, K., Dobbins, T., Dawson, A., Isbister, G., & Buckley, N. (2017). Relationship between prescribed psychotropic medications and co-ingested alcohol in intentional self-poisonings. *British Journal of Psychiatry*, 210: 203-208 .
- Newton-Howes, G., Foulds, J., Guy, N., Boden, J., & Mulder, R. (2017). Personality disorder and alcohol treatment outcome: systematic review and meta-analysis. *The British Journal of Psychiatry*, 211:22-30.

'555' Topics (5 slides on each topic with no more than 5 bullet points)

- Personality disorder and substance misuse
- Depression and alcohol
- Psychotropic drug interactions with opioid substitution medications
- Public health concerns of Chemsex

MCQs

1. Which of the following have not been associated with misuse potential:

- A. Acamprosate
- B. Hyoscine butylbromide
- C. Loperamide
- D. Pregabalin
- E. Codeine phosphate

2. Intoxication of the following substances can be associated with psychosis in DSM 5 except :

- A. Alcohol
- B. Cannabis
- C. Sedatives
- D. Opioids
- E. Inhalants

3. Which of the following symptoms is not associated with Benzodiazepines withdrawal using CIWA B

- A. Loss of appetite
- B. Yawning
- C. Problems sleeping
- D. Difficulties with concentration

E. Sensitivity to light / blurred vision

4. Percentage of patients attending Community Mental Health Teams reporting past-year problem drug use and/or harmful alcohol use has been found to be approximately:

- A. 24%
- B. 34%
- C. 44%
- D. 54%
- E. 64%

5. Which of the following is likely to predate Alcohol and other Drug use disorder in most cases;

- A. Generalised anxiety disorder (GAD),
- B. Panic disorder,
- C. Depression
- D. Dysthymia
- E. PTSD

EMI Questions

Drugs that may induce psychiatric symptoms:

- A. Gamma-Hydroxybutyric acid (GHB)
- B. Lysergic acid diethylamide (LSD)
- C. Ketamine
- D. Phencyclidine (PCP)
- E. Diazepam
- F. Amphetamine
- G. Cocaine
- H. Alcohol
- I. Cannabis
- J. Butane

1a. This psychoactive component of this drug acts through the type 1 form of the receptors which are found in high concentrations throughout the cerebellum, hippocampus, basal ganglia, cortex, brainstem, thalamus and hypothalamus

1b. This compound acts as an agonist at 5HT_{2A} receptor

1c. One of the main mechanisms of action of this drug is by reverse transfer of the neurotransmitter dopamine

Psychotropic medications used in people with co-occurring mental health and alcohol/drug use conditions:

- A. Diazepam
- B. Quetiapine
- C. Risperidone
- D. Citalopram
- E. Amisulpride
- F. Sertraline
- G. Baclofen
- H. Olanzapine
- I. Aripipazole
- J. Fluoxetine

2a. Disulfiram can inhibit the metabolism of this compound

2b. This antipsychotic should be considered in patients with impaired liver function

2c. This agent may increase risk of compulsive behaviour such as gambling

MCQ Answers

Q1 A Acamprosate (Used for alcohol dependence)

Q2 D Opioids see DSM 5

Q3 B Yawning in COWS

Q4 C 44% see Weaver (2003)

Q5 E PTSD see Swendsen (2010)

EMI 1

(1) I (Nestler 2008)

(2) B

(3) F Cocaine works mainly by reuptake inhibition

EMI 2

(1) A

(2) E

(3) I

Additional Resources / Reading Materials

E-Learning

Royal College of Psychiatrists CPD Online

- Dual diagnosis: the diagnosis and treatment of depression with co-existing substance misuse.

Journal Articles

- Agabio, R., Trogu, E., & Pani, P. (2018, 4). Antidepressants for the treatment of people with co-occurring depression and alcohol dependence. *The Cochrane database of systematic reviews*, 4, CD008581.
- Aichhorn, W., Santeler, S., Stelzig-Schöler, R., Kemmler, G., Steinmayr-Gensluckner, M., & Hinterhuber, H. (2008). Prevalence of psychiatric disorders among homeless adolescents. *Neuropsychiatrie: Klinik, Diagnostik, Therapie und Rehabilitation: Organ der Gesellschaft Österreichischer Nervenärzte und Psychiater*, 22(3), 180-188.
- Baandrup L, Ebdrup BH, Rasmussen JØ, Lindschou J, Gluud C, Glenthøj BY. Pharmacological interventions for benzodiazepine discontinuation in chronic benzodiazepine users. *Cochrane Database of Systematic Reviews* 2018, Issue 3. Art. No.: CD011481. DOI: 10.1002/14651858.CD011481.pub2.
- Bebbington, P., & McManus, S. (2009). Adult psychiatric morbidity in England, 2007 : results of a household survey. London: National Centre for Social Research.
- Beijer, U., Andréasson, A., Ågren, G., & Fugelstad, A. (2007). Mortality, mental disorders and addiction: a 5-year follow-up of 82 homeless men in Stockholm. *Nordic Journal of Psychiatry*, 61(5), 363-368.
- Busto, U.E., Sykora, K. & Sellers, E.M. (1989). A clinical scale to assess benzodiazepine withdrawal. *Journal of Clinical Psychopharmacology*, 9 (6), 412–416.
- Caton, C., Hasin, D., Shrout, P., Drake, R., Dominguez, B., Samet, S., & Schanzer, B. (2006). Predictors of psychosis remission in psychotic disorders that co-occur with substance use. *Schizophrenia Bulletin*, 32(4), 618-25.
- Clinical Guidelines on Drug Misuse and Dependence Update 2017 Independent Expert Working Group (2017). Drug misuse and dependence: UK guidelines on clinical management. London: Department of Health.
- Colizzi, M., & Murray, R. (2018, 4 20). Cannabis and psychosis: what do we know and what should we do? *The British Journal of Psychiatry*, 212(04), 195-196.
- Combaluzier, S., B. Gouvernet, and A. Bernoussi. "Impact of personality disorders in a sample of 212 homeless drug users." *L'Encephale* 35.5 (2009): 448-453.

- Conner, K. R., Piquart, M., & Duberstein, P. R. (2008). Meta-analysis of depression and substance use and impairment among intravenous drug users (IDUs). *Addiction*, 103(4), 524-534.
- Conner, K. R., Piquart, M., & Gamble, S. A. (2009). Meta-analysis of depression and substance use among individuals with alcohol use disorders. *Journal of Substance Abuse Treatment*, 37(2), 127-137.
- Conner, K. R., Piquart, M., & Holbrook, A. P. (2008). Meta-analysis of depression and substance use and impairment among cocaine users. *Drug and Alcohol Dependence*, 98(1-2), 13-23.
- Crump, C., Sundquist, K., Winkleby, M. A., & Sundquist, J. (2013). Comorbidities and mortality in bipolar disorder: a Swedish national cohort study. *JAMA psychiatry*, 70(9), 931-939.
- Crunelle, C. L., Van Den Brink, W., Moggi, F., Konstenius, M., Franck, J., Levin, F. R., ... & Schellekens, A. (2018). International consensus statement on screening, diagnosis and treatment of substance use disorder patients with comorbid attention Deficit/Hyperactivity disorder. *European addiction research*, 24(1), 43-51.
- Darker CD, Sweeney BP, Barry JM, Farrell MF, Donnelly-Swift E. Psychosocial interventions for benzodiazepine harmful use, abuse or dependence. *Cochrane Database of Systematic Reviews* 2015, Issue 5. Art. No.: CD009652. DOI: 10.1002/14651858.CD009652.pub2.
- Davis, L. L., Pilkinton, P., Wisniewski, S. R., Trivedi, M. H., Gaynes, B. N., Howland, R. H., et al. (2012). Effect of concurrent substance use disorder on the effectiveness of single and combination antidepressant medications for the treatment of major depression: an exploratory analysis of a single-blind randomized trial. *Depression and anxiety*, 29(2), 111-122.
- Delgadillo, J. G., C. Gilbody, S. Payne, S. (2013). Depression, anxiety and comorbid substance use: association patterns in outpatient addictions treatment. *Mental Health and Substance Use*, 6(1), 59-75.
- Demyttenaere, Koen, et al. "Prevalence, severity, and unmet need for treatment of mental disorders in the World Health Organization World Mental Health Surveys." *Jama* 291.21 (2004): 2581-2590
- Haw, C. M., & Hawton, K. (2011). Problem drug use, drug misuse and deliberate self-harm: trends and patient characteristics, with a focus on young people, Oxford, 1993–2006. *Social psychiatry and psychiatric epidemiology*, 46(2), 85-93.
- Dimeff, L. A., & Linehan, M. M. (2008). Dialectical behavior therapy for substance abusers. *Addiction science & clinical practice*, 4(2), 39.

- Farnia, V., Shakeri, J., Tatari, F., Juibari, T. A., Yazdchi, K., Bajoghli, H., ... & Aghaei, A. (2014). Randomized controlled trial of aripiprazole versus risperidone for the treatment of amphetamine-induced psychosis. *The American journal of drug and alcohol abuse*, 40(1), 10-15.
- Fazel, S., Smith, E. N., Chang, Z., & Geddes, J. R. (2018). Risk factors for interpersonal violence: an umbrella review of meta-analyses. *The British Journal of Psychiatry*, 213(4), 609-614.
- Foulds, J., Adamson, S., Boden, J., Williman, J., & Mulder, R. (2015). Depression in patients with alcohol use disorders: Systematic review and meta-analysis of outcomes for independent and substance-induced disorders. *Journal of Affective Disorders*, 185:47-59.
- Gibbon S, Duggan C, Stoffers J, Huband N, Völlm BA, Ferriter M, Lieb K. Psychological interventions for antisocial personality disorder. Cochrane Database of Systematic Reviews 2010, Issue 6. Art. No.: CD007668. DOI: 10.1002/14651858.CD007668.pub2.
- Gimeno, C., Dorado, M. L., Roncero, C., Szerman, N., Vega, P., Balanzá-Martínez, V., & Alvarez, F. J. (2017). Treatment of comorbid alcohol dependence and anxiety disorder: review of the scientific evidence and recommendations for treatment. *Frontiers in psychiatry*, 8, 1730.
- Gold, A. K., Otto, M. W., Deckersbach, T., Sylvia, L. G., Nierenberg, A. A., & Kinrys, G. (2018). Substance use comorbidity in bipolar disorder: A qualitative review of treatment strategies and outcomes. *The American journal on addictions*, 27(3), 188-201.
- Goldstein, R. B., Smith, S. M., Chou, S. P., Saha, T. D., Jung, J., Zhang, H., ... & Grant, B. F. (2016). The epidemiology of DSM-5 posttraumatic stress disorder in the United States: results from the National Epidemiologic Survey on Alcohol and Related Conditions-III. *Social psychiatry and psychiatric epidemiology*, 51(8), 1137-1148.
- Healthcare Quality Improvement Partnership. (2018). National Confidential Inquiry into Suicide and Homicide: Report 2018.
- Hunt, G. E., Siegfried, N., Morley, K., Sitharthan, T., & Cleary, M. (2013). Psychosocial interventions for people with both severe mental illness and substance misuse. *Cochrane Database of Systematic Reviews*, (10).
- Hides, L., Samet, S., & Lubman, D. I. (2010). Cognitive behaviour therapy (CBT) for the treatment of co-occurring depression and substance use: Current evidence and directions for future research. *Drug and Alcohol Review*, 29(5), 508-517.
- Hoskins, Mathew, et al. "Pharmacotherapy for post-traumatic stress disorder: systematic review and meta-analysis." *The British Journal of Psychiatry* 206.2 (2015): 93-100.
- Iovieno, N., Tedeschini, E., Bentley, K., Evins, a., & Papakostas, G. (2011). Antidepressants for major depressive disorder and dysthymic disorder in patients with

comorbid alcohol use disorders: a meta-analysis of placebo-controlled randomized trials. *The Journal of clinical psychiatry*, 72 (8), 1144-51.

- Kushner, M. G., Maurer, E. W., Thuras, P., Donahue, C., Frye, B., Menary, K. R., ... & Van Demark, J. (2013). Hybrid cognitive behavioral therapy versus relaxation training for co-occurring anxiety and alcohol disorder: A randomized clinical trial. *Journal of consulting and clinical psychology*, 81(3), 429.
- Lee, N. K., Jenner, L., Harney, A., & Cameron, J. (2018). Pharmacotherapy for amphetamine dependence: A systematic review. *Drug and alcohol dependence*.
- Marel, C., Mills, K. L., Kingston, R., Gournay, K., Deady, M., Kay-Lambkin, F., ... & Teesson, M. (2016). *Co-occurring alcohol and other drug and mental health conditions in alcohol and other drug treatment settings*. Illustrations.
- Maremmani, A. G., Rovai, L., Rugani, F., Bacciardi, S., Dell'osso, L., & Maremmani, I. (2014). Substance abuse and psychosis. The strange case of opioids. *Eur Rev Med Pharmacol Sci*, 18(3), 287-302.
- Martinez-Raga, J., Knecht, C., De Alvaro, R., Szerman, N., & Ruiz, P. (2013). Addressing dual diagnosis patients suffering from attention-deficit hyperactivity disorders and comorbid substance use disorders: A review of treatment considerations. *Addictive Disorders & Their Treatment*, 12(4), 213-230.
- Martín-Santos, R., Fonseca, F., Domingo-Salvany, A., Ginés, J. M., Ímaz, M. L., Navinés, R., ... & Torrens, M. (2006). Dual diagnosis in the psychiatric emergency room in Spain. *The European Journal of Psychiatry*, 20(3), 147-156.
- McHugh, R. Kathryn. "Treatment of co-occurring anxiety disorders and substance use disorders." *Harvard review of psychiatry* 23.2 (2015): 99.
- Messer, Thomas, et al. "Substance abuse in patients with bipolar disorder: A systematic review and meta-analysis." *Psychiatry research* 253 (2017): 338-350.
- Messer, T., Lammers, G., Müller-Siecheneder, F., Schmidt, R. F., & Latifi, S. (2017). Substance abuse in patients with bipolar disorder: A systematic review and meta-analysis. *Psychiatry research*, 253, 338-350.
- Messina, N., Farabee, D., & Rawson, R. (2003). Treatment responsivity of cocaine-dependent patients with antisocial personality disorder to cognitive-behavioral and contingency management interventions. *Journal of consulting and clinical psychology*, 71(2), 320.
- Mestre-Pintó, J. I., Domingo-Salvany, A., Martín-Santos, R., & Torrens, M. (2014). Dual diagnosis screening interview to identify psychiatric comorbidity in substance users: development and validation of a brief instrument. *European addiction research*, 20(1), 41-48.

- National Institute for Health and Care Excellence. (2018). Attention deficit hyperactivity disorder: diagnosis and management. London: National Institute for Health and Care Excellence.
- National Institute for Health and Care Excellence (2018). Post-traumatic stress disorder NG 116. (2018). London: National Institute for Health and Care Excellence.
- National Institute for Health and Care Excellence. (2011). Psychosis with coexisting substance misuse CG120. London: National Institute for Health and Care Excellence.
- Niemi-Pynttäre, J., Sund, R., Putkonen, H., Vormaa, H., Wahlbeck, K., & Pirkola, S. (2013). Substance-induced psychoses converting into schizophrenia: A register-based study of 18,478 Finnish inpatient cases. *Journal of Clinical Psychiatry*, 74(1), e94-9.
- Nunes E V, Levin F R. (2004) Treatment of depression in patients with alcohol or other drug dependence: a meta-analysis. *JAMA*, 291(15), 1887-1896.
- Petrakis, I. L., Poling, J., Levinson, C., Nich, C., Carroll, K., Ralevski, E., & Rounsaville, B. (2006). Naltrexone and disulfiram in patients with alcohol dependence and comorbid post-traumatic stress disorder. *Biological Psychiatry*, 60(7), 777-783.
- Pettinati, H., O'Brien, C., & Dundon, W. (2013). Current status of co-occurring mood and substance use disorders: A new therapeutic target. *American Journal of Psychiatry*, 170(1), 23–30.
- PHE. (2017). Better care for people with co-occurring mental health, and alcohol and drug use conditions.
- Radhakrishnan, R., Wilkinson, S. T., & D'Souza, D. C. (2014). Gone to Pot - A Review of the Association between Cannabis and Psychosis. *Front Psychiatry*, 5, 54.
- Riper, H., Andersson, G., Hunter, S., de Wit, J., Berking, M., & Cuijpers, P. (2014). Treatment of comorbid alcohol use disorders and depression with cognitive-behavioural therapy and motivational interviewing: A meta-analysis. *Addiction*, 109(3), 394-406.
- Roberts, N. P., Roberts, P. A., Jones, N., & Bisson, J. I. (2015). Psychological interventions for post-traumatic stress disorder and comorbid substance use disorder: A systematic review and meta-analysis. *Clinical psychology review*, 38, 25-38.
- Sepede, G., Lorusso, M., Spano, M. C., Di Nanno, P., Di Iorio, G., & Di Giannantonio, M. (2018). Efficacy and Safety of Atypical Antipsychotics in Bipolar Disorder With Comorbid Substance Dependence: A Systematic Review. *Clinical neuropharmacology*, 41(5), 181-191.
- Shoptaw, S. J., Kao, U., & Ling, W. W. (2008). Treatment for amphetamine psychosis. *Cochrane database of systematic reviews*, (4).

- Starzer, M., Nordentoft, M., & Hjorthøj, C. (2018). Rates and predictors of conversion to schizophrenia or bipolar disorder following substance-induced psychosis. *American Journal of Psychiatry*, 175(4), 343-350.
- Swendsen, J., Conway, K. P., Degenhardt, L., Glantz, M., Jin, R., Merikangas, K. R., ... & Kessler, R. C. (2010). Mental disorders as risk factors for substance use, abuse and dependence: results from the 10-year follow-up of the National Comorbidity Survey. *Addiction*, 105(6), 1117-1128.
- Thomsen, K. R., Thylstrup, B., Pedersen, M. M., Pedersen, M. U., Simonsen, E., & Hesse, M. (2018). Drug-related predictors of readmission for schizophrenia among patients admitted to treatment for drug use disorders. *Schizophrenia research*, 195, 495-500.
- Tiihonen, J., Mittendorfer-Rutz, E., Torniainen, M., Alexanderson, K., & Tanskanen, A. (2015). Mortality and cumulative exposure to antipsychotics, antidepressants, and benzodiazepines in patients with schizophrenia: an observational follow-up study. *American Journal of Psychiatry*, 173(6), 600-606.
- Tirado-Muñoz, J., Farré, A., Mestre-Pintó, J., Szerman, N., & Torrens, M. (2018). Dual diagnosis in Depression: treatment recommendations. *adicciones*, 30(1).
- Torrens, M., Mestre-Pintó, J. I., & Domingo-Salvany, A. (2015). *Comorbidity of substance use and mental disorders in Europe*. Publication Office of the European Union.
- Van de Glind, G., van den Brink, W., Koeter, M. W., Carpentier, P. J., van Emmerik-van Oortmerssen, K., Kaye, S., ... & Moggi, F. (2013). Validity of the Adult ADHD Self-Report Scale (ASRS) as a screener for adult ADHD in treatment seeking substance use disorder patients. *Drug and alcohol dependence*, 132(3), 587-596.
- van Emmerik-van Oortmerssen, K., van de Glind, G., Koeter, M. W., Allsop, S., Auriacombe, M., Barta, C., ... & Casas, M. (2014). Psychiatric comorbidity in treatment-seeking substance use disorder patients with and without attention deficit hyperactivity disorder: results of the IASP study. *Addiction*, 109(2), 262-272.
- van Emmerik-van Oortmerssen, K., van de Glind, G., van den Brink, W., Smit, F., Crunelle, C. L., Swets, M., & Schoevers, R. A. (2012). Prevalence of attention-deficit hyperactivity disorder in substance use disorder patients: a meta-analysis and meta-regression analysis. *Drug and alcohol dependence*, 122(1-2), 11-19.
- van Emmerik-van Oortmerssen, K., Vedel, E., Kramer, F. J., Blankers, M., Dekker, J. J., van den Brink, W., & Schoevers, R. A. (2019). Integrated cognitive behavioral therapy for ADHD in adult substance use disorder patients: results of a randomized clinical trial. *Drug and alcohol dependence*, 197, 28-36.

- Volkow, N. D., Swanson, J. M., Evins, A. E., DeLisi, L. E., Meier, M. H., Gonzalez, R., ... & Baler, R. (2016). Effects of cannabis use on human behavior, including cognition, motivation, and psychosis: a review. *JAMA psychiatry*, 73(3), 292-297.
- Weaver, T., Madden, P., Charles, V., Stimson, G., Renton, A., Tyrer, P., et al. (2003). Comorbidity of substance misuse and mental illness in community mental health and substance misuse services. *The British Journal of Psychiatry*, 183, 304-313.
- Weiss, R. D., Griffin, M. L., Kolodziej, M. E., Greenfield, S. F., Najavits, L. M., Daley, D. C., ... & Hennen, J. A. (2007). A randomized trial of integrated group therapy versus group drug counseling for patients with bipolar disorder and substance dependence. *American Journal of Psychiatry*, 164(1), 100-107.
- Xue, X., Song, Y., Yu, X., Fan, Q., Tang, J., & Chen, X. (2018). Olanzapine and haloperidol for the treatment of acute symptoms of mental disorders induced by amphetamine-type stimulants: A randomized controlled trial. *Medicine*, 97(8).
- Zarse, E. M., Neff, M. R., Yoder, R., Hulvershorn, L., Chambers, J. E., & Chambers, R. A. (2019). The adverse childhood experiences questionnaire: two decades of research on childhood trauma as a primary cause of adult mental illness, addiction, and medical diseases. *Cogent Medicine*, 6(1), 1581447.